

# **Product datasheet for MC204216**

## Sfrp1 (NM\_013834) Mouse Untagged Clone

**Product data:** 

**Product Type:** Expression Plasmids

**Product Name:** Sfrp1 (NM\_013834) Mouse Untagged Clone

Tag: Tag Free Symbol: Sfrp1

**Synonyms:** 2210415K03Rik; AW011917; AW107218; AW742929; sFRP-1

Mammalian Cell

Selection:

Neomycin

Vector: PCMV6-Kan/Neo (PCMV6KN)

E. coli Selection: Kanamycin (25 ug/mL)

Fully Sequenced ORF: >BC024495

GCAGCCCGCAGCCCGCCGCCCCTCTGCGAGCCGGGACAGCACTCGGCCCCGCGCGCTCCCCGCCCCGCGC CGCCGAGCCGGACGGGGCCCGGCACTGCGCCTTTGTCCCCGGAGGCTCCGGGAAGTTTGCAGCGGGACGC GCGCGTGAAGGCAGCGTGGGCAGCCCCGACGTCGCCGAGCACATGGGCGTCGGGCGCAGCGCGCGGGGGT CGCGGCGGGGCCGCCTCGGGAGTGCTGCTGGCGTTGGCCGCCGCTCTGCTGGCCTCGGGTTCGGCCAGCG AGTACGACTACGTGAGCTTCCAGTCCGACATCGGCTCGTATCAGAGCGGGCGCTTCTACACCAAGCCCCC GCAGTGCGTGGACATCCCGGTGGACCTGAGGCTGTGCCACAACGTGGGCTACAAGAAGATGGTGCTGCCC CTACCCGTGTCGCTGGCTCTGCGAGGCCGTGCGCGACTCGTGCGAGCCGGTCATGCAGTTCTTCGGCTTC TACTGGCCCGAGATGCTCAAATGTGACAAGTTCCCCGAGGGCGACGTCTGCATCGCCATGACCCCGCCCA ATACCACGGAAGCCTCTAAGCCCCAAGGTACAACCGTGTGTCCTCCATGCGACAACGAGTTGAAGTCAGA GGCCATCATTGAACATCTCTGTGCAAGCGAGTTTGCACTGAGGATGAAAATCAAAGAAGTGAAGAAGGAA AACGGTGACAAGAAGATTGTCCCCAAGAAGAAGAACCCTTGAAGCTGGGGCCCATCAAGAAGAAGGAGC TGAAGCGGCTTGTGCTGTTCCTGAAGAACGGTGCCGACTGTCCCTGCCACCAGCTGGACAACCTCAGCCA CAACTTTCTCATCATGGGCCGCAAGGTGAAGAGCCAGTACCTGCTGACAGCCATTCACAAGTGGGACAAG AAAAACAAGGAGTTCAAAAACTTCATGAAGAGAATGAAAAACCACGAGTGTCCCACCTTCCAGTCTGTTT TTAAGTGATACTGGGGCGGACTGGGGAAGGGGAGTGTGGCTTGGGGTGAGGGTGGGGCGCGTGGATGAC

**Restriction Sites:** Rsrll-Notl

**ACCN:** NM\_013834

**Insert Size:** 945 bp



**OriGene Technologies, Inc.** 9620 Medical Center Drive, Ste 200

CN: techsupport@origene.cn

Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com



#### **OTI Disclaimer:**

Due to the inherent nature of this plasmid, standard methods to replicate additional amounts of DNA in E. coli are highly likely to result in mutations and/or rearrangements. Therefore, OriGene does not guarantee the capability to replicate this plasmid DNA. Additional amounts of DNA can be purchased from OriGene with batch-specific, full-sequence verification at a reduced cost. Please contact our customer care team at <a href="mailto:customport@origene.com">customport@origene.com</a> or by calling 301.340.3188 option 3 for pricing and delivery.

The molecular sequence of this clone aligns with the gene accession number as a point of reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing variants is recommended prior to use. <u>More info</u>

### Components:

The ORF clone is ion-exchange column purified and shipped in a 2D barcoded Matrix tube containing 10ug of transfection-ready, dried plasmid DNA (reconstitute with 100 ul of water).

### **Reconstitution Method:**

- 1. Centrifuge at 5,000xg for 5min.
- 2. Carefully open the tube and add 100ul of sterile water to dissolve the DNA.
- 3. Close the tube and incubate for 10 minutes at room temperature.
- 4. Briefly vortex the tube and then do a quick spin (less than 5000xg) to concentrate the liquid at the bottom.
- 5. Store the suspended plasmid at -20°C. The DNA is stable for at least one year from date of shipping when stored at -20°C.

Note:

Plasmids are not sterile. For experiments where strict sterility is required, filtration with 0.22um filter is required.

**RefSeq:** <u>BC024495</u>, <u>AAH24495</u>

 RefSeq Size:
 1340 bp

 RefSeq ORF:
 945 bp

 Locus ID:
 20377

 UniProt ID:
 Q8C4U3

 Cytogenetics:
 8 11.48 cM

**Gene Summary:** 

Soluble frizzled-related proteins (sFRPS) function as modulators of Wnt signaling through direct interaction with Wnts. They have a role in regulating cell growth and differentiation in specific cell types. SFRP1 decreases intracellular beta-catenin levels (By similarity). Has antiproliferative effects on vascular cells, in vitro and in vivo, and can induce, in vivo, an angiogenic response. In vascular cell cycle, delays the G1 phase and entry into the S phase. In kidney development, inhibits tubule formation and bud growth in metanephroi (By similarity). Inhibits WNT1/WNT4-mediated TCF-dependent transcription (By similarity). [UniProtKB/Swiss-Prot Function]