

Product datasheet for **MC204692**

Snrpg (NM_026506) Mouse Untagged Clone

Product data:

Product Type:	Expression Plasmids
Product Name:	Snrpg (NM_026506) Mouse Untagged Clone
Tag:	Tag Free
Symbol:	Snrpg
Synonyms:	2810024K17Rik; AL022803; sm-G; SMG; snRNP-G
Mammalian Cell Selection:	Neomycin
Vector:	PCMV6-Kan/Neo (PCMV6KN)
E. coli Selection:	Kanamycin (25 ug/mL)
Fully Sequenced ORF:	>BC027499 CCGCAGAGGAGTCGCCATGAGCAAAGCCCACCCTCCCGAGCTGAAGAAGTTTATGGACAAGAAGTTATCA TTGAAGTTAAACGGTGGCAGGCATGTCCAAGGAATACTGCGGGGCTTTGATCCCTTTATGAACCTTGTA TTGACGAGTGTGTGGAGATGGCAACCAAGTGGGCAACAGAACAACATTGGCATGGTGGTCATCCGAGGAAA CAGCATCATCATGTTAGAAGCCTTGAAAAGAGTCTGACCTGTGCTCAGCAAGCAGTGTCCACATCCCTCC CCAAAGGCCTGTTTGATTGTGATGTAGAATTAGGTGATGACATTTTCATATGGAACCTTTTACTAAATA AACTTTTGTGATACTCAAAAAAAAAAAAAAAAAAAAAAAAAAAAAAAAAAAAAA
Restriction Sites:	EcoRI-NotI
ACCN:	NM_026506
Insert Size:	231 bp
OTI Disclaimer:	Our molecular clone sequence data has been matched to the reference identifier above as a point of reference. Note that the complete sequence of our molecular clones may differ from the sequence published for this corresponding reference, e.g., by representing an alternative RNA splicing form or single nucleotide polymorphism (SNP).
Components:	The ORF clone is ion-exchange column purified and shipped in a 2D barcoded Matrix tube containing 10ug of transfection-ready, dried plasmid DNA (reconstitute with 100 ul of water).



[View online »](#)

Reconstitution Method:

1. Centrifuge at 5,000xg for 5min.
2. Carefully open the tube and add 100ul of sterile water to dissolve the DNA.
3. Close the tube and incubate for 10 minutes at room temperature.
4. Briefly vortex the tube and then do a quick spin (less than 5000xg) to concentrate the liquid at the bottom.
5. Store the suspended plasmid at -20°C. The DNA is stable for at least one year from date of shipping when stored at -20°C.

RefSeq: [BC027499](#), [AAH27499](#)

RefSeq Size: 398 bp

RefSeq ORF: 231 bp

Locus ID: 68011

UniProt ID: [P62309](#)

Cytogenetics: 6 D1

Gene Summary: Plays role in pre-mRNA splicing as core component of the SMN-Sm complex that mediates spliceosomal snRNP assembly and as component of the spliceosomal U1, U2, U4 and U5 small nuclear ribonucleoproteins (snRNPs), the building blocks of the spliceosome. Component of both the pre-catalytic spliceosome B complex and activated spliceosome C complexes. Is also a component of the minor U12 spliceosome. As part of the U7 snRNP it is involved in histone 3'-end processing.[UniProtKB/Swiss-Prot Function]