

## **Product datasheet for MC204897**

## Nudt5 (NM\_016918) Mouse Untagged Clone

**Product data:** 

**Product Type:** Expression Plasmids

Product Name: Nudt5 (NM\_016918) Mouse Untagged Clone

Tag: Tag Free
Symbol: Nudt5

Mammalian Cell Neomycin

Selection:

Vector:

PCMV6-Kan/Neo (PCMV6KN)

E. coli Selection: Kanamycin (25 ug/mL)

Fully Sequenced ORF: >BC049595

Restriction Sites: Sfil-Sfil

**ACCN:** NM\_016918

**Insert Size:** 657 bp

**OTI Disclaimer:** Our molecular clone sequence data has been matched to the reference identifier above as a

point of reference. Note that the complete sequence of our molecular clones may differ from the sequence published for this corresponding reference, e.g., by representing an alternative

RNA splicing form or single nucleotide polymorphism (SNP).



**OriGene Technologies, Inc.** 9620 Medical Center Drive, Ste 200

CN: techsupport@origene.cn

Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com



**Components:** The ORF clone is ion-exchange column purified and shipped in a 2D barcoded Matrix tube

containing 10ug of transfection-ready, dried plasmid DNA (reconstitute with 100 ul of water).

**Reconstitution Method:** 1. Centrifuge at 5,000xg for 5min.

2. Carefully open the tube and add 100ul of sterile water to dissolve the DNA.

3. Close the tube and incubate for 10 minutes at room temperature.

4. Briefly vortex the tube and then do a quick spin (less than 5000xg) to concentrate the liquid

at the bottom.

5. Store the suspended plasmid at -20°C. The DNA is stable for at least one year from date of

shipping when stored at -20°C.

**RefSeq:** <u>BC049595</u>, <u>AAH49595</u>

RefSeq Size: 1053 bp
RefSeq ORF: 657 bp
Locus ID: 53893
UniProt ID: Q9|KX6
Cytogenetics: 2 A1

**Gene Summary:** Enzyme that can either act as an ADP-sugar pyrophosphatase in absence of diphosphate or

catalyze the synthesis of ATP in presence of diphosphate (By similarity). In absence of diphosphate, hydrolyzes with similar activities various modified nucleoside diphosphates

such as ADP-ribose, ADP-mannose, ADP-glucose, 8-oxo-GDP and 8-oxo-dGDP (PubMed:10722730). Can also hydrolyze other nucleotide sugars with low activity (PubMed:10722730). In presence of diphosphate, mediates the synthesis of ATP in the nucleus by catalyzing the conversion of ADP-ribose to ATP and ribose 5-phosphate (By

similarity). Nuclear ATP synthesis takes place when dephosphorylated at Thr-44 (By similarity). Nuclear ATP generation is required for extensive chromatin remodeling events that are

(PubMed:21070968). Binds U8 snoRNA (PubMed:21070968).[UniProtKB/Swiss-Prot Function]

energy-consuming (By similarity). Does not play a role in U8 snoRNA decapping activity