

Product datasheet for MC204921

Ndufa6 (NM_025987) Mouse Untagged Clone

Product data:

Product Type: Expression Plasmids

Product Name: Ndufa6 (NM_025987) Mouse Untagged Clone

Tag: Tag Free Symbol: Ndufa6

Synonyms: 14kDa; 2700038D15Rik; B230217P19Rik

Mammalian Cell

Selection:

Neomycin

Vector: PCMV6-Kan/Neo (PCMV6KN)

E. coli Selection: Kanamycin (25 ug/mL)

Fully Sequenced ORF: >BC048451

Restriction Sites: Sfil-Sfil

ACCN: NM_025987

Insert Size: 396 bp

OTI Disclaimer: Our molecular clone sequence data has been matched to the reference identifier above as a

point of reference. Note that the complete sequence of our molecular clones may differ from the sequence published for this corresponding reference, e.g., by representing an alternative

RNA splicing form or single nucleotide polymorphism (SNP).

Components: The ORF clone is ion-exchange column purified and shipped in a 2D barcoded Matrix tube

containing 10ug of transfection-ready, dried plasmid DNA (reconstitute with 100 ul of water).



OriGene Technologies, Inc. 9620 Medical Center Drive, Ste 200

CN: techsupport@origene.cn

Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com



Reconstitution Method:

- 1. Centrifuge at 5,000xg for 5min.
- 2. Carefully open the tube and add 100ul of sterile water to dissolve the DNA.
- 3. Close the tube and incubate for 10 minutes at room temperature.
- 4. Briefly vortex the tube and then do a quick spin (less than 5000xg) to concentrate the liquid at the bottom.

5. Store the suspended plasmid at -20°C. The DNA is stable for at least one year from date of shipping when stored at -20°C.

RefSeq: <u>BC048451</u>, <u>AAH48451</u>

RefSeq Size: 567 bp
RefSeq ORF: 396 bp
Locus ID: 67130
UniProt ID: Q9CQZ5
Cytogenetics: 15 E1

Gene Summary: Accessory subunit of the mitochondrial membrane respiratory chain NADH dehydrogenase

(Complex I), that is believed to be not involved in catalysis. Required for proper complex I assembly. Complex I functions in the transfer of electrons from NADH to the respiratory chain. The immediate electron acceptor for the enzyme is believed to be ubiquinone.

[UniProtKB/Swiss-Prot Function]