

Product datasheet for **MC205684**

Smarcd3 (NM_025891) Mouse Untagged Clone

Product data:

Product Type: Expression Plasmids
Product Name: Smarcd3 (NM_025891) Mouse Untagged Clone
Tag: Tag Free
Symbol: Smarcd3
Synonyms: 1500001J14Rik; 2210409C08Rik; BAF60C
Mammalian Cell Selection: Neomycin
Vector: PCMV6-Kan/Neo (PCMV6KN)
E. coli Selection: Kanamycin (25 ug/mL)
Fully Sequenced ORF: >BC013122

```

CTTTTTTCTCCAACCTCTCCCCTCTGAGTCCTGCTGGGCCCTCCCGGTCCACTCGAGTAGAAGTG
TGAGAGAGCCCATCGGGACTCAGTGGGAAGGCTGGGTTTATTCAAAGGCCGAAAAGGGAAGAAAGAGG
AAGAGAGAGAGAGAGTGTAGAGGAGCCGCTGAGCCACCCCGATGGCCGCGGACGAAAGTTGCCGAGGGGC
GCGCAAAGCCACGAAAAGCAAACCTTTTTGAGTTTCTGGTCCATGGGGTGCGCCCGGGATGCCGTCTGGA
GCCCGAATGCCACCAGGGGGCGCCATGGGCCCGGGCTCCCGTACATGGGCAGCCCCGGGTAC
GACCCGGCTGGCCCCGCGGGCATGGAGCCCGCCGCAAGCGAGCAGCGCCCCCGGGCAGAGCCA
GGCACAGGGCCAGGGCCAGCCCTGCCACCGCCAGCGGGAGCCGAGTCCCAAGAGGAGGAAGATG
GCTGACAAAATCTCCCTCAAAGGATTCGGGAGCTGGTACCCGAGTCCAGGCTTACATGGACCTCCTAG
CATTTGAGAGGAACTGGATCAAACCATCATGCGGAAGCGGGTGGACATCCAGGAGCCCTGAAGAGGCC
CATGAAGCAAAGCGAAAGCTGCGCCTTTATATCTCCAATACTTTTAAACCCTGCGAAGCCTGATGCGGAA
GACTCTGATGGCAGCATTGCCTCCTGGGAGCTGCGGGTGGAGGGGAAGCTTTGGATGATCCTAGTAAGC
AGAAGAGGAAGTTTTCTTCTTCTCAAGAGTTTGGTCATTGAGTTGGACAAAGACCTTTATGGCCAGA
CAACCACCTTGTGAGTGGCACCAGGACCCACAACCCAGGAAACAGATGGGTTCCAAGTGAAGAGACCA
GGGACTTGAGTGTGCGCTGCACCCTGCTCCTGATGCTGGACTATCAGCCTCCCCAGTTCAAATTGGACC
CCCGCTTAGCCCGGCTGCTGGGGTTACACACACAGAGCCGCTCAGCCATTGTCCAGGCACTGTGGCAGTA
TGTGAAGACCAACAGGCTACAGGACTCCCATGACAAGGAGTACATCAATGGCGACAAGTATTTCCAGCAG
ATTTTTGACTGCCCCGCTAAAGTTCTCTGAGATTCCCAGCGCCTCACAGCCCTGCTGCTGCCCTG
ACCCATTGTGATCAACCACGTATCAGCGTGGACCCATCAGACCAGAAGAAGACAGCGTGTATGACAT
AGATGTGGAGGTGGAGGAACCGCTGAAAGGGCAGATGAGTAGCTTCTCCTGTCCACGGCAACCCAGCAG
GAGATCAGTGCTCTGGACAGTAAGATCCATGAGACGATTGAGTCCATAAACCCAGCTCAAGATCCAGAGGG
ACTTCATGCTAAGTTTCTCCAGAGACCCAAAGGCTACGTCCAAGACCTGCTCCGCTCCCAGAGCCGTGA
TCTCAAGGTGATGACAGATGTGGCAGGGAACCCCGAGGAAGAACGCAGGGCTGAGTTCTACCACCAGCCC
TGGTCCCAGGAAGCCGTTAGCCGCTACTTCTACTGTAAGATCCAGCAGCGCAGGCAGGAGCTGGAGCAGT
CGCTGGTCTGCGCAACACCTAGGAGCCCGTGAACAAGCGTCAGGGTGGACCAGCCACTCCGCCAGCAC
AGGCCCTGGGCTCTGACTCCCCCTCTCGCGCTGTGCGGAAGGTGGGAGGGCTGGATGGATTAAGGTC
ACGTAACAGACAAAAA

```



[View online »](#)

Restriction Sites:	RsrII-NotI
ACCN:	NM_025891
Insert Size:	1452 bp
OTI Disclaimer:	Our molecular clone sequence data has been matched to the reference identifier above as a point of reference. Note that the complete sequence of our molecular clones may differ from the sequence published for this corresponding reference, e.g., by representing an alternative RNA splicing form or single nucleotide polymorphism (SNP).
Components:	The ORF clone is ion-exchange column purified and shipped in a 2D barcoded Matrix tube containing 10ug of transfection-ready, dried plasmid DNA (reconstitute with 100 ul of water).
Reconstitution Method:	<ol style="list-style-type: none">1. Centrifuge at 5,000xg for 5min.2. Carefully open the tube and add 100ul of sterile water to dissolve the DNA.3. Close the tube and incubate for 10 minutes at room temperature.4. Briefly vortex the tube and then do a quick spin (less than 5000xg) to concentrate the liquid at the bottom.5. Store the suspended plasmid at -20°C. The DNA is stable for at least one year from date of shipping when stored at -20°C.
RefSeq:	BC013122 , AAH13122
RefSeq Size:	1790 bp
RefSeq ORF:	1452 bp
Locus ID:	66993
UniProt ID:	Q6P9Z1
Cytogenetics:	5 A3
Gene Summary:	Involved in transcriptional activation and repression of select genes by chromatin remodeling (alteration of DNA-nucleosome topology). Component of SWI/SNF chromatin remodeling complexes that carry out key enzymatic activities, changing chromatin structure by altering DNA-histone contacts within a nucleosome in an ATP-dependent manner (PubMed:22952240, PubMed:26601204). Stimulates nuclear receptor mediated transcription. Belongs to the neural progenitors-specific chromatin remodeling complex (npBAF complex) and the neuron-specific chromatin remodeling complex (nBAF complex). During neural development a switch from a stem/progenitor to a postmitotic chromatin remodeling mechanism occurs as neurons exit the cell cycle and become committed to their adult state. The transition from proliferating neural stem/progenitor cells to postmitotic neurons requires a switch in subunit composition of the npBAF and nBAF complexes. As neural progenitors exit mitosis and differentiate into neurons, npBAF complexes which contain ACTL6A/BAF53A and PHF10/BAF45A, are exchanged for homologous alternative ACTL6B/BAF53B and DPF1/BAF45B or DPF3/BAF45C subunits in neuron-specific complexes (nBAF). The npBAF complex is essential for the self-renewal/proliferative capacity of the multipotent neural stem cells. The nBAF complex along with CREST plays a role regulating the activity of genes essential for dendrite growth (PubMed:17640523).[UniProtKB/Swiss-Prot Function]