

Product datasheet for **MC206553**

Atox1 (BC027632) Mouse Untagged Clone

Product data:

Product Type:	Expression Plasmids
Product Name:	Atox1 (BC027632) Mouse Untagged Clone
Tag:	Tag Free
Symbol:	Atox1
Synonyms:	AI256639; Atx1
Mammalian Cell Selection:	Neomycin
Vector:	PCMV6-Kan/Neo (PCMV6KN)
E. coli Selection:	Kanamycin (25 ug/mL)
Fully Sequenced ORF:	>BC027632 CAGTCGTCGTCGCCGCTGCCACTGCCCGTGTGTGCCGCGTCAGTCATGCCGAAGCAGGATTCTCCGTGG ACATGACCTGTGAGGGCTGTGCTGAAGCCGTCTCCAGAGTCCTCAACAAGCTGGGAGGAGTGGAGTTCAA CATTGACCTGCCCAACAAGAAGGTCTGCATCGACTCTGAGCACAGCTCAGACACCCTGCTGGCAACCCTC AACAAAACAGGAAAGGCTGTTTCTACCTTGGCCCAAGTAGCCAGGACCTGGGCGAGTCCCTCCGGATA TAAACTGAAGAGGCAGGCTGTTGATCTGGTCTCCCGGCAGATCTGGAACACCAACTGCTCAGTCCAGTC CAGCCCAGCCATGGAGTTCCTGCCAGACAGGCCCTCCCGCTGGCTCCCTGCAAGCTTCCATGTAATAA AGTCAAGCTGAGTTTGTAAA AAAAAAAAAA
Restriction Sites:	EcoRI-NotI
ACCN:	BC027632
Insert Size:	207 bp
OTI Disclaimer:	Our molecular clone sequence data has been matched to the reference identifier above as a point of reference. Note that the complete sequence of our molecular clones may differ from the sequence published for this corresponding reference, e.g., by representing an alternative RNA splicing form or single nucleotide polymorphism (SNP).
Components:	The ORF clone is ion-exchange column purified and shipped in a 2D barcoded Matrix tube containing 10ug of transfection-ready, dried plasmid DNA (reconstitute with 100 ul of water).



[View online »](#)

Reconstitution Method:

1. Centrifuge at 5,000xg for 5min.
2. Carefully open the tube and add 100ul of sterile water to dissolve the DNA.
3. Close the tube and incubate for 10 minutes at room temperature.
4. Briefly vortex the tube and then do a quick spin (less than 5000xg) to concentrate the liquid at the bottom.
5. Store the suspended plasmid at -20°C. The DNA is stable for at least one year from date of shipping when stored at -20°C.

RefSeq: [BC027632](#), [AAH27632](#)

RefSeq Size: 501 bp

RefSeq ORF: 207 bp

Locus ID: 11927

Cytogenetics: 11 33.07 cM

Gene Summary: Binds and deliver cytosolic copper to the copper ATPase proteins. May be important in cellular antioxidant defense (By similarity).[UniProtKB/Swiss-Prot Function]