

Product datasheet for **MC206717**

Ap2s1 (BC052499) Mouse Untagged Clone

Product data:

Product Type: Expression Plasmids
Product Name: Ap2s1 (BC052499) Mouse Untagged Clone
Tag: Tag Free
Symbol: Ap2s1
Synonyms: MGC62945
Mammalian Cell Selection: Neomycin
Vector: PCMV6-Kan/Neo (PCMV6KN)
E. coli Selection: Kanamycin (25 ug/mL)
Fully Sequenced ORF: >BC052499
 GGCACGAGGCACAACCTGCAAGCAGAGCCGAGCCGCCCTGTTACAGCGGGACCCGGGCTCTGGGGTCCA
 GACAGGGGTCGCCATGATCCGATTCATCCTTATCCAGTGGTACATGCAGTTCGATGACGACGAGAAGCAG
 AAGCTGATCGAGGAGGTGCACGCCGTGGTCACCGTCAGGGATGCCAAGCACACCAACTTTGTGGAGTTCC
 GGAACTTCAAGATCATCTACCGACGCTACGCTGGCCTCTACTTCTGCATCTGCGTGGATGTCAACGACAA
 CAATCTGGCCTATCTCGAGGCCATCCACAACCTTCGTAGAAGTGTTAAATGAATACTTCCACAATGTCTGT
 GAACTGGACCTGGTGTCAACTTCTACAAGTTTACACGGTGGTAGATGAGATGTTCTGGCAGGAGAGA
 TCCGAGAGACCAGCCAGACGAAGGTGCTGAAGCAGCTGCTGATGCTGCAGTCCCTGGAGTGAGGGGGCT
 CTGCCCAGCCCGCCTGCTGGACCAACCTGCCTGCTGCTCCCCTGCCTGCCAGGCCTGAGGCCACCCTA
 GCAGCCCCCTTCCCTTCTCAGCTGGCCCAGAGGAAGGACCCGGCAGGGTCTAGGCCACAGGGGAGGAGGC
 CAGAACCCACTGCCTCTGGGCCCTCCGTGGATGGCAGAGGCCACCGTGATGTCCCAGTAAGTGTGCCAT
 TGTCGTGTGATGCTGTGAGTGTGTGTGTGTGTAGCCCCCAATAAACCTGTGGTCTGCCTGAAAAA
 AAAGAA

Restriction Sites: EcoRI-NotI

ACCN: BC052499

Insert Size: 399 bp

OTI Disclaimer: Our molecular clone sequence data has been matched to the reference identifier above as a point of reference. Note that the complete sequence of our molecular clones may differ from the sequence published for this corresponding reference, e.g., by representing an alternative RNA splicing form or single nucleotide polymorphism (SNP).

Components: The ORF clone is ion-exchange column purified and shipped in a 2D barcoded Matrix tube containing 10ug of transfection-ready, dried plasmid DNA (reconstitute with 100 ul of water).


[View online »](#)

Reconstitution Method:

1. Centrifuge at 5,000xg for 5min.
2. Carefully open the tube and add 100ul of sterile water to dissolve the DNA.
3. Close the tube and incubate for 10 minutes at room temperature.
4. Briefly vortex the tube and then do a quick spin (less than 5000xg) to concentrate the liquid at the bottom.
5. Store the suspended plasmid at -20°C. The DNA is stable for at least one year from date of shipping when stored at -20°C.

Note: Plasmids are not sterile. For experiments where strict sterility is required, filtration with 0.22um filter is required.

RefSeq: [BC052499](#), [AAH52499](#)

RefSeq Size: 829 bp

RefSeq ORF: 399 bp

Locus ID: 232910

Cytogenetics: 7 A2

Gene Summary: Component of the adaptor protein complex 2 (AP-2). Adaptor protein complexes function in protein transport via Transport vesicles in different membrane traffic pathways. Adaptor protein complexes are vesicle coat components and appear to be involved in cargo selection and vesicle formation. AP-2 is involved in clathrin-dependent endocytosis in which cargo proteins are incorporated into vesicles surrounded by clathrin (clathrin-coated vesicles, CCVs) which are destined for fusion with the early endosome. The clathrin lattice serves as a mechanical scaffold but is itself unable to bind directly to membrane components. Clathrin-associated adaptor protein (AP) complexes which can bind directly to both the clathrin lattice and to the lipid and protein components of membranes are considered to be the major clathrin adaptors contributing the CCV formation. AP-2 also serves as a cargo receptor to selectively sort the membrane proteins involved in receptor-mediated endocytosis. AP-2 seems to play a role in the recycling of synaptic vesicle membranes from the presynaptic surface. AP-2 recognizes Y-X-X-[FILMV] (Y-X-X-Phi) and [ED]-X-X-X-L-[LI] endocytosis signal motifs within the cytosolic tails of transmembrane cargo molecules. AP-2 may also play a role in maintaining normal post-endocytic trafficking through the ARF6-regulated, non-clathrin pathway. The AP-2 alpha and AP-2 sigma subunits are thought to contribute to the recognition of the [ED]-X-X-X-L-[LI] motif. May also play a role in extracellular calcium homeostasis (By similarity).[UniProtKB/Swiss-Prot Function]