

Product datasheet for **MC206925**

Hbxip (BC028547) Mouse Untagged Clone

Product data:

Product Type:	Expression Plasmids
Product Name:	Hbxip (BC028547) Mouse Untagged Clone
Tag:	Tag Free
Symbol:	Hbxip
Synonyms:	1110003H18Rik; Hbxip; XIP
Mammalian Cell Selection:	Neomycin
Vector:	pCMV6-Entry (PS100001)
E. coli Selection:	Kanamycin (25 ug/mL)
Fully Sequenced ORF:	>MC206925 representing BC028547. Blue=ORF Red=Cloning site Green=Tag(s)

GCTCGTTTAGTGAACCGTCAGAATTTGTAAACGACTCACTATAGGGCGCCGGGAATTCGTCGACTG
GATCCGGTACCGAGGAGATCTGCCGCC**GCGATCGCC**
ATGGAGGCGACTTTGGAGCAGCATTGGAGGACACAATGAAGAATCCATCCATTGTTGGAGTCTATGC
ACAGATTCACAAGGACTTAATCTGGGCTGCCGTGGTACCCTGTCGGATGAGCACGCTGGAGTCATATCT
GTTCTAGCCAGCAGGCAGCTAGGCTAACCTCTGACCCACCGACATCCCTGTGGTATGTTAGAATCA
GATAATGGGAACATTATGATCCAGAAACACGATGGCATCACAGTGGCTGTGCACAAAATGGCCTCTGA


ACGCGTACGCGGCCGCTCGAGCAGAAACTCATCTCAGAAGAGGATCTGGCAGCAAATGATATCCTGGAT
TACAAGGATGACGACGATAAGGTTTAAACGGCCGGC

Restriction Sites: Sgfl-Mlul



[View online »](#)

Plasmid Map:



ACCN: BC028547

Insert Size: 276 bp

OTI Disclaimer: Our molecular clone sequence data has been matched to the reference identifier above as a point of reference. Note that the complete sequence of our molecular clones may differ from the sequence published for this corresponding reference, e.g., by representing an alternative RNA splicing form or single nucleotide polymorphism (SNP).

Components: The ORF clone is ion-exchange column purified and shipped in a 2D barcoded Matrix tube containing 10ug of transfection-ready, dried plasmid DNA (reconstitute with 100 ul of water).

Reconstitution Method:

1. Centrifuge at 5,000xg for 5min.
2. Carefully open the tube and add 100ul of sterile water to dissolve the DNA.
3. Close the tube and incubate for 10 minutes at room temperature.
4. Briefly vortex the tube and then do a quick spin (less than 5000xg) to concentrate the liquid at the bottom.
5. Store the suspended plasmid at -20°C. The DNA is stable for at least one year from date of shipping when stored at -20°C.

RefSeq: [BC028547](#)

RefSeq Size: 735 bp

RefSeq ORF: 275 bp

Locus ID: 68576

Cytogenetics: 3 F2.3

MW: 9.6 kDa

Gene Summary: As part of the Ragulator complex it is involved in amino acid sensing and activation of mTORC1, a signaling complex promoting cell growth in response to growth factors, energy levels, and amino acids. Activated by amino acids through a mechanism involving the lysosomal V-ATPase, the Ragulator functions as a guanine nucleotide exchange factor activating the small GTPases Rag. Activated Ragulator and Rag GTPases function as a scaffold recruiting mTORC1 to lysosomes where it is in turn activated. When complexed to BIRC5, interferes with apoptosome assembly, preventing recruitment of pro-caspase-9 to oligomerized APAF1, thereby selectively suppressing apoptosis initiated via the mitochondrial/cytochrome c pathway (By similarity).[UniProtKB/Swiss-Prot Function]