

## Product datasheet for **MC207106**

### 0610008C08Rik (BC016557) Mouse Untagged Clone

#### Product data:

**Product Type:** Expression Plasmids  
**Product Name:** 0610008C08Rik (BC016557) Mouse Untagged Clone  
**Tag:** Tag Free  
**Symbol:** 0610008C08Rik  
**Synonyms:** 0610008C08Rik; 1110019O03Rik  
**Mammalian Cell Selection:** Neomycin  
**Vector:** pCMV6-Entry (PS100001)  
**E. coli Selection:** Kanamycin (25 ug/mL)  
**Fully Sequenced ORF:** >BC016557  
 Red=Cloning site Blue=ORF Green=Tags(s)

TTTTGTAATACGACTCACTATAGGGCGCCGGGAATTCGTCGACTGGATCCGGTACCGAGGAGATCTGCC  
 GCC**CGATCGCC**

ATGAAGATCGATGAGCTTTCCTACTCTACTCAGTTCCTGAGGGTCAATCTAAATATGTGGAGGAGCCAAGGA  
 CTCAACTGAAGAAAACATCTCACAACCTCCGACATCATTGTGAGCCATATACAAGTTTCTGTCAGGAAAT  
 AACTCCCATACTAAACCAAGGTGGATCACTTGTCCAGTGGGAGTAGACAACATAACTATCTTCAA  
 AATGCGCCTCCTGGATTTTCCCAAGACTCGGTGTTATTGGTTTTGCTGGTTTTGTTGGACTCCTTTTTG  
 CTAGAGGTTCAAAAATAAAGAAGCTGGTGTATCCTCCTTTTTTTCATGGGATTAGGTGCCTCTGTCTATTA  
 CCCACAACAAGCCATCACCATTGCCAGATCACTGGGGAGAAGTTATATGACTGGGGATTACGAGGGTAC  
 ATAGTTATAGAAGATTTGTGGAAGCAAAATTTTCAGAAGCCAGGAAATGTGAAGAATTCACCTGGAATA  
 AATAG

**ACGCGT**ACGCGGCCGCTCGAGCAGAACTCATCTCTGAAGAGGATCTGGCAGCAAATGATATCCTGGATT  
 ACAAGGATGACGACGATAAGGTTTAA

**Restriction Sites:** Sgfl-Mlul  
**ACCN:** BC016557  
**Insert Size:** 495 bp

**OTI Disclaimer:** Our molecular clone sequence data has been matched to the reference identifier above as a point of reference. Note that the complete sequence of our molecular clones may differ from the sequence published for this corresponding reference, e.g., by representing an alternative RNA splicing form or single nucleotide polymorphism (SNP).



[View online »](#)

<b>Components:</b>	The ORF clone is ion-exchange column purified and shipped in a 2D barcoded Matrix tube containing 10ug of transfection-ready, dried plasmid DNA (reconstitute with 100 ul of water).
<b>Reconstitution Method:</b>	<ol style="list-style-type: none"><li>1. Centrifuge at 5,000xg for 5min.</li><li>2. Carefully open the tube and add 100ul of sterile water to dissolve the DNA.</li><li>3. Close the tube and incubate for 10 minutes at room temperature.</li><li>4. Briefly vortex the tube and then do a quick spin (less than 5000xg) to concentrate the liquid at the bottom.</li><li>5. Store the suspended plasmid at -20°C. The DNA is stable for at least one year from date of shipping when stored at -20°C.</li></ol>
<b>Note:</b>	Plasmids are not sterile. For experiments where strict sterility is required, filtration with 0.22um filter is required.
<b>RefSeq:</b>	<a href="#">BC016557</a> , <a href="#">AAH16557</a>
<b>RefSeq Size:</b>	1186 bp
<b>RefSeq ORF:</b>	494 bp
<b>Locus ID:</b>	68316
<b>Cytogenetics:</b>	X C3
<b>Gene Summary:</b>	Component of the MICOS complex, a large protein complex of the mitochondrial inner membrane that plays crucial roles in the maintenance of crista junctions, inner membrane architecture, and formation of contact sites to the outer membrane. Plays a crucial role in crista junction formation and mitochondrial function (By similarity). Can induce cardiac lipotoxicity by enhancing mitochondrial respiration and fatty acid metabolism in cardiac myoblasts (PubMed:24743151). Promotes cholesterol efflux from macrophage cells. Detected in HDL, LDL and VLDL. Secreted by a microsomal triglyceride transfer protein (MTTP)-dependent mechanism, probably as a VLDL-associated protein that is subsequently transferred to HDL (By similarity).[UniProtKB/Swiss-Prot Function]