

Product datasheet for **MC207128**

2900042B11Rik (BC021897) Mouse Untagged Clone

Product data:

Product Type: Expression Plasmids
Product Name: 2900042B11Rik (BC021897) Mouse Untagged Clone
Tag: Tag Free
Symbol: 2900042B11Rik
Synonyms: 2900042B11Rik; IFT25
Vector: pCMV6-Entry (PS100001)
E. coli Selection: Kanamycin (25 ug/mL)
Cell Selection: Neomycin
Fully Sequenced ORF: >BC021897
Red=Cloning site Blue=ORF Green=Tags(s)

TTTTGTAATACGACTCACTATAGGGCGGCCGGAATTCGTCGACTGGATCCGGTACCGAGGAGATCTGCC
GCC**GCGATCGCC**

ATGAGGAAAGTGGATCTCTGCTCAGTCACTGAAGGGACAGAAGTGATTCTGGCCACATCGAGTGATGAAA
AGCACCCACCTGAAAACATCATTGATGGTCCTGCTTGGCGAGGACGTAGAGGGAAGTCGTATTCTCAGAC
CTTTCATTCCAAGAAGAGTGGCCAATGGAGAACACACCACCTTTTGTATTCTAATTTTATTTATAT
GTGGGTTTTGCCTACATGTATGGCTGTGCGGATATGTTCACTACCTAG

ACGCGTACGCGGCCGCTCGAGCAGAACTCATCTCTGAAGAGGATCTGGCAGCAAATGATATCCTGGATT
ACAAGGATGACGACGATAAGGTTTAA

Restriction Sites: Sgfl-Mlul
ACCN: BC021897
Insert Size: 258 bp

OTI Disclaimer: Our molecular clone sequence data has been matched to the reference identifier above as a point of reference. Note that the complete sequence of our molecular clones may differ from the sequence published for this corresponding reference, e.g., by representing an alternative RNA splicing form or single nucleotide polymorphism (SNP).

Components: The ORF clone is ion-exchange column purified and shipped in a 2D barcoded Matrix tube containing 10ug of transfection-ready, dried plasmid DNA (reconstitute with 100 ul of water).



[View online »](#)

Reconstitution Method:

1. Centrifuge at 5,000xg for 5min.
2. Carefully open the tube and add 100ul of sterile water to dissolve the DNA.
3. Close the tube and incubate for 10 minutes at room temperature.
4. Briefly vortex the tube and then do a quick spin (less than 5000xg) to concentrate the liquid at the bottom.
5. Store the suspended plasmid at -20°C. The DNA is stable for at least one year from date of shipping when stored at -20°C.

RefSeq: [BC021897](#), [AAH21897](#)

RefSeq Size: 1537 bp

RefSeq ORF: 257 bp

Locus ID: 72938

Cytogenetics: 4 C7

Gene Summary: Component of the IFT complex B required for sonic hedgehog/SHH signaling. May mediate transport of SHH components: required for the export of SMO and PTCH1 receptors out of the cilium and the accumulation of GLI2 at the ciliary tip in response to activation of the SHH pathway, suggesting it is involved in the dynamic transport of SHH signaling molecules within the cilium. Not required for ciliary assembly (PubMed:22595669). Its role in intraflagellar transport is mainly seen in tissues rich in ciliated cells such as kidney and testis. Essential for male fertility, spermiogenesis and sperm flagella formation (PubMed:28430876). Plays a role in the early development of the kidney (PubMed:29626631). May be involved in the regulation of ureteric bud initiation (PubMed:29626631).[UniProtKB/Swiss-Prot Function]