

Product datasheet for MC207134

Bloc1s2 (NM_028607) Mouse Untagged Clone

Product data:

Product Type: Expression Plasmids

Product Name: Bloc1s2 (NM_028607) Mouse Untagged Clone

Tag: Tag Free
Symbol: Bloc1s2

Synonyms: 2410089B13Rik; Bloc1s2a; BLOS2

Mammalian Cell

Selection:

Neomycin

Vector:pCMV6-Entry (PS100001)E. coli Selection:Kanamycin (25 ug/mL)

Fully Sequenced ORF: >MC207134 representing NM_028607

Red=Cloning site Blue=ORF Orange=Stop codon

TTTTGTAATACGACTCACTATAGGGCGGCCGGGAATTCGTCGACTGGATCCGGTACCGAGGAGATCTGCC

GCCGCGATCGCC

GAGAAGCGATGA

ACGCGTACGCGGCCGCTCGAGCAGAAACTCATCTCAGAAGAGGATCTGGCAGCAAATGATATCCTGGATT

ACAAGGATGACGACGATAAGGTTTAA

Restriction Sites: Sgfl-Mlul
ACCN: NM_028607
Insert Size: 432 bp

OTI Disclaimer: Our molecular clone sequence data has been matched to the reference identifier above as a

point of reference. Note that the complete sequence of our molecular clones may differ from the sequence published for this corresponding reference, e.g., by representing an alternative

RNA splicing form or single nucleotide polymorphism (SNP).



OriGene Technologies, Inc. 9620 Medical Center Drive, Ste 200

CN: techsupport@origene.cn

Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com



Components: The O

The ORF clone is ion-exchange column purified and shipped in a 2D barcoded Matrix tube containing 10ug of transfection-ready, dried plasmid DNA (reconstitute with 100 ul of water).

Reconstitution Method:

- 1. Centrifuge at 5,000xg for 5min.
- 2. Carefully open the tube and add 100ul of sterile water to dissolve the DNA.
- 3. Close the tube and incubate for 10 minutes at room temperature.
- 4. Briefly vortex the tube and then do a quick spin (less than 5000xg) to concentrate the liquid

at the bottom.

5. Store the suspended plasmid at -20°C. The DNA is stable for at least one year from date of

shipping when stored at -20°C.

RefSeq: <u>NM 028607.1</u>, <u>NP 082883.1</u>

RefSeq Size: 910 bp
RefSeq ORF: 432 bp
Locus ID: 73689
UniProt ID: Q9CWG9
Cytogenetics: 19 C3

Gene Summary: Component of the BLOC-1 complex, a complex that is required for normal biogenesis of

lysosome-related organelles (LRO), such as platelet dense granules and melanosomes (By similarity). In concert with the AP-3 complex, the BLOC-1 complex is required to target

membrane protein cargos into vesicles assembled at cell bodies for delivery into neurites and nerve terminals (PubMed:16760431, PubMed:21998198). The BLOC-1 complex, in association

with SNARE proteins, is also proposed to be involved in neurite extension

(PubMed:19546860). As part of the BORC complex may play a role in lysosomes movement and localization at the cell periphery. Associated with the cytosolic face of lysosomes, the BORC complex may recruit ARL8B and couple lysosomes to microtubule plus-end-directed

kinesin motor (By similarity).[UniProtKB/Swiss-Prot Function]