

Product datasheet for MC207174

OriGene Technologies, Inc. 9620 Medical Center Drive, Ste 200

Rockville, MD 20850, US
Phone: +1-888-267-4436
https://www.origene.com
techsupport@origene.com
EU: info-de@origene.com
CN: techsupport@origene.cn

Sfrs8 (BC034336) Mouse Untagged Clone

Product data:

Product Type: Expression Plasmids

Product Name: Sfrs8 (BC034336) Mouse Untagged Clone

Tag: Tag Free Symbol: Sfrs8

Synonyms: 1190005N23Rik; 6330437E22Rik; Al197402; AW212079; Sfrs8; Srsf8; SWAP

Vector:pCMV6-Entry (PS100001)E. coli Selection:Kanamycin (25 ug/mL)

Cell Selection: Neomycin

Fully Sequenced ORF: >MC207174 representing BC034336

Red=Cloning site Blue=ORF

TTTTGTAATACGACTCACTATAGGGCGGCCGGGAATTCGTCGACTGGATCCGGTACCGAGGAGATCTGCC

GCCGCGATCGCC

ACGCGTACGCGGCCGCTCGAGCAGAAACTCATCTCAGAAGAGGATCTGGCAGCAAATGATATCCTGGATT

ACAAGGATGACGACGATAAGGTTTAA

Restriction Sites: Sgfl-Mlul
ACCN: BC034336
Insert Size: 267 bp

OTI Disclaimer: Our molecular clone sequence data has been matched to the reference identifier above as a

point of reference. Note that the complete sequence of our molecular clones may differ from the sequence published for this corresponding reference, e.g., by representing an alternative

RNA splicing form or single nucleotide polymorphism (SNP).

Components: The ORF clone is ion-exchange column purified and shipped in a 2D barcoded Matrix tube

containing 10ug of transfection-ready, dried plasmid DNA (reconstitute with 100 ul of water).







Reconstitution Method:

- 1. Centrifuge at 5,000xg for 5min.
- 2. Carefully open the tube and add 100ul of sterile water to dissolve the DNA.
- 3. Close the tube and incubate for 10 minutes at room temperature.
- 4. Briefly vortex the tube and then do a quick spin (less than 5000xg) to concentrate the liquid

at the bottom.

5. Store the suspended plasmid at -20°C. The DNA is stable for at least one year from date of

shipping when stored at -20°C.

RefSeq: <u>BC034336.1</u>

RefSeq Size: 1806 bp
RefSeq ORF: 266 bp
Locus ID: 231769
Cytogenetics: 5 G1.3

Gene Summary: Plays a role as an alternative splicing regulator. Regulates its own expression at the level of

RNA processing. Also regulates the splicing of fibronectin and CD45 genes. May act, at least in

part, by interaction with other R/S-containing splicing factors. Represses the splicing of

MAPT/Tau exon 10 (By similarity).[UniProtKB/Swiss-Prot Function]