

Product datasheet for **MC207176**

Gtf2a2 (BC028551) Mouse Untagged Clone

Product data:

Product Type: Expression Plasmids
Product Name: Gtf2a2 (BC028551) Mouse Untagged Clone
Tag: Tag Free
Symbol: Gtf2a2
Synonyms: 12kDa, Tfllg
Vector: pCMV6-Entry (PS100001)
E. coli Selection: Kanamycin (25 ug/mL)
Cell Selection: Neomycin
Fully Sequenced ORF: >BC028551
Red=Cloning site Blue=ORF Green=Tags(s)

TTTTGTAATACGACTCACTATAGGGCGGCCGGAATTCGTCGACTGGATCCGGTACCGAGGAGATCTGCC
GCC**GCGATCGCC**

**ATGGCCTATCAGTTATACAGAAATACAACCTTTGGGGAACAGTCTTCAAGAGAGCCTTGATGAGCTCATA
AGTCTCAACAGATCACCCCCAGCTTGCCTTCAAGTTCTACTTCAGTTTGATAAAGCTATAAATTCAGC
ATTGGCTCAGAGAGTCAGGAACAGAGTCAATTCAGGAAAATCTGGGTTTGGTCCAGCACCTATGTCA
GGTGGTCACAACCTACCCGTGA**

**ACGCGTACGCGGCCGCTCGAGCAGAAACTCATCTCTGAAGAGGATCTGGCAGCAAATGATATCCTGGATT
ACAAGGATGACGACGATAAGGTTTAA**

Restriction Sites: Sgfl-Mlul
ACCN: BC028551
Insert Size: 231 bp

OTI Disclaimer: Our molecular clone sequence data has been matched to the reference identifier above as a point of reference. Note that the complete sequence of our molecular clones may differ from the sequence published for this corresponding reference, e.g., by representing an alternative RNA splicing form or single nucleotide polymorphism (SNP).

Components: The ORF clone is ion-exchange column purified and shipped in a 2D barcoded Matrix tube containing 10ug of transfection-ready, dried plasmid DNA (reconstitute with 100 ul of water).



[View online »](#)

Reconstitution Method:

1. Centrifuge at 5,000xg for 5min.
2. Carefully open the tube and add 100ul of sterile water to dissolve the DNA.
3. Close the tube and incubate for 10 minutes at room temperature.
4. Briefly vortex the tube and then do a quick spin (less than 5000xg) to concentrate the liquid at the bottom.
5. Store the suspended plasmid at -20°C. The DNA is stable for at least one year from date of shipping when stored at -20°C.

RefSeq: [BC028551](#), [AAH28551](#)

RefSeq Size: 919 bp

RefSeq ORF: 230 bp

Locus ID: 235459

Cytogenetics: 9 D

Gene Summary: TFIIA is a component of the transcription machinery of RNA polymerase II and plays an important role in transcriptional activation. TFIIA in a complex with TBP mediates transcriptional activity (By similarity).[UniProtKB/Swiss-Prot Function]