

## Product datasheet for **MC207544**

### Maea (NM\_021500) Mouse Untagged Clone

#### Product data:

Product Type: Expression Plasmids  
 Product Name: Maea (NM\_021500) Mouse Untagged Clone  
 Tag: Tag Free  
 Symbol: Maea  
 Synonyms: 1110030D19Rik; EMP; Gid9  
 Vector: pCMV6-Entry (PS100001)  
 E. coli Selection: Kanamycin (25 ug/mL)  
 Cell Selection: Neomycin  
 Fully Sequenced ORF: >MC207544 representing NM\_021500  
 Red=Cloning site Blue=ORF Orange=Stop codon

TTTTGTAATACGACTCACTATAGGGCGCCGGAATTCGTCGACTGGATCCGGTACCGAGGAGATCTGCC  
 GCC**CGATCGCC**

ATGGCGGTGCAGGAGTCGGCGGCCAGCTGTCCATGACCCTGAAGGTGCAGGAGTACCCGACCCTCAAGG  
 TGCCCTATGAGACTGAACAAACGCTTCCGAGCTGCTCAGAAAAACATCGATCGAGAGACTAGCCACGT  
 CACCATGGTGGTAGCTGAGCTTGAAGAAGACCTTGAGTAGTTGCCAGCTGTGGACTCTGTGGTCAGCCTA  
 TTGGATGGTGTGGTGGAGAAGCTGAGTGTCTCAAGAGGAAGGCAGTAGAGTCCATCCAGGCCGAGGATG  
 AGAGCGCCAAGCTCTGCAACCTAGGATCGAGCACCTCAAGGAGCACAGCAGTGACCAGCCAGCAGCAGC  
 CAGCATGTGGAAGCGGAAGCGCATGGACCGGATGATGGTGGAGCATCTGCTACGCTGTGGCTACTACAAC  
 ACAGCTGTGAAGCTGGCTCGCCAGAGTGGCATCGAGGACCTTGTGAATATCGAGATGTTCTGACAGCCA  
 AAGAAGTGGAGGAGTCTTGGAGAGGCGTGAGACAGCCACCTGCCTTGCCTGGTCCATGATAACAAGTC  
 CCGACTCCGGAAGATGAAGAGCTGCCTAGAGTTCAGCCTCAGGATTCAGGAGTTCATTGAACTTGTCCGG  
 CAGAACAAGCGCCTGGATGCTGTGAGACATGCAAGAAAGCACTTCAGTCAGGCTGAAGGGAGCCAGCTGG  
 ATGAGGTCCGCCAGGTCATGGGCATGTTGGCCTTCCCACCAGACACATATCTCCCATACAAGGACCT  
 CCTGGACCCAGCCGGTGGCAATGCTGATCCAGCAGTTTCGATATGATAACTACCGGCTGCACCAGCTG  
 GGAAACAGCTCAGTCTTACCCTCACCTGCAGGCTGGGCTCTCAGCAATAAAGACACCACAGTGCTACA  
 AGGAGGATGGCAGCTCTAAGAGCCCTGACTGCCCTGTGTGCAGCCGCTCTCTGAACAAACTGGCAGCC  
 CCTGCCCATGGCTCACTGTGCCAACTCCCCTGGTCTGCAAGATCTCTGGTGACGTGATGAATGAGAAT  
 AACCCACCATGATGCTGCCTAATGGCTATGTCTATGGCTACAATTCTCTGCTTTCTATTCGTCGAAGATG  
 ATAAAGTTGTTTGCCCAAGAACCAAGAAGTCTCCACTTCTCCAAGCTGAGAAAGTATACATCATGTA  
 G

**ACGCGT**ACGCGGCCGCTCGAGCAGAAACTCATCTCAGAAGAGGATCTGGCAGCAAATGATATCCTGGATT  
 ACAAGGATGACGACGATAAGGTTTAA



[View online »](#)

<b>Restriction Sites:</b>	Sgfl-Mlul
<b>ACCN:</b>	NM_021500
<b>Insert Size:</b>	1191 bp
<b>OTI Disclaimer:</b>	Our molecular clone sequence data has been matched to the reference identifier above as a point of reference. Note that the complete sequence of our molecular clones may differ from the sequence published for this corresponding reference, e.g., by representing an alternative RNA splicing form or single nucleotide polymorphism (SNP).
<b>Components:</b>	The ORF clone is ion-exchange column purified and shipped in a 2D barcoded Matrix tube containing 10ug of transfection-ready, dried plasmid DNA (reconstitute with 100 ul of water).
<b>Reconstitution Method:</b>	<ol style="list-style-type: none"><li>1. Centrifuge at 5,000xg for 5min.</li><li>2. Carefully open the tube and add 100ul of sterile water to dissolve the DNA.</li><li>3. Close the tube and incubate for 10 minutes at room temperature.</li><li>4. Briefly vortex the tube and then do a quick spin (less than 5000xg) to concentrate the liquid at the bottom.</li><li>5. Store the suspended plasmid at -20°C. The DNA is stable for at least one year from date of shipping when stored at -20°C.</li></ol>
<b>RefSeq:</b>	<u><a href="#">NM_021500.2</a></u> , <u><a href="#">NP_067475.2</a></u>
<b>RefSeq Size:</b>	2128 bp
<b>RefSeq ORF:</b>	1191 bp
<b>Locus ID:</b>	59003
<b>UniProt ID:</b>	<u><a href="#">Q4VC33</a></u>
<b>Cytogenetics:</b>	5 B1
<b>Gene Summary:</b>	Core component of the CTLH E3 ubiquitin-protein ligase complex that selectively accepts ubiquitin from UBE2H and mediates ubiquitination and subsequent proteasomal degradation of the transcription factor HBP1. MAEA and RMND5A are both required for catalytic activity of the CTLH E3 ubiquitin-protein ligase complex. MAEA is required for normal cell proliferation. The CTLH E3 ubiquitin-protein ligase complex is not required for the degradation of enzymes involved in gluconeogenesis, such as FBP1 (By similarity). Plays a role in erythroblast enucleation during erythrocyte maturation and in the development of mature macrophages (PubMed:16707498). Mediates the attachment of erythroid cell to mature macrophages; this MAEA-mediated contact inhibits erythroid cell apoptosis (By similarity). Participates in erythroblastic island formation, which is the functional unit of definitive erythropoiesis (PubMed:16707498, PubMed:17071116). Associates with F-actin to regulate actin distribution in erythroblasts and macrophages (PubMed:16707498). May contribute to nuclear architecture and cells division events (By similarity).[UniProtKB/Swiss-Prot Function]