

Product datasheet for MC207675

Chmp2a (NM_026885) Mouse Untagged Clone

Product data:

| | |
|---------------------------|---------------------------------------------------------------------------------|
| Product Type: | Expression Plasmids |
| Product Name: | Chmp2a (NM_026885) Mouse Untagged Clone |
| Tag: | Tag Free |
| Symbol: | Chmp2a |
| Synonyms: | 1500016L11Rik; mVps2 |
| Mammalian Cell Selection: | Neomycin |
| Vector: | pCMV6-Entry (PS100001) |
| E. coli Selection: | Kanamycin (25 ug/mL) |
| Fully Sequenced ORF: | >MC207675 representing NM_026885 Red=Cloning site Blue=ORF Orange=Stop codon |

TTTGTAAATACGACTCACTATAGGGCGGCCGGAATTCGTCGACTGGATCCGGTACCGAGGAGATCTGCC
 GCCCGCATCGCC

ATGGACCTGTTGTTTGGGCGCCGGAAGACGCCAGAGGAACTACTTCGGCAAAACCAGAGGGCCCTGAACC
 GAGCCATGAGAGAACTGGACAGGGAACGACAGAACTAGAAACCCAGGAAAAGAAAATCATTGCGGACAT
 CAAAAAGATGGCAAAGCAAGGCCAGATGGATGCTGTTTGAATCATGGCAAAAGACCTGGTGCCTACCCGG
 AGATATGTACGCAAGTTTGTGTTGATGCGGGCCAACTCAAGCTGTGTCCCTCAAGATACAGACTCTAA
 AATCCAACAACCTCAATGGCACAAGCCATGAAGGGTGTACTAAGGCCATGGGCACTATGAACAGACAGCT
 GAAATTACCCAGATCCAGAAGATCATGATGGAGTTTGAACGGCAGGCAGAGATCATGGACATGAAGGAA
 GAAATGATGAATGACGAATTGATGATGCCATGGGTGATGAGGAAGATGAAGAGGAGAGTGATGCTGTTG
 TGTCCCAGGTCCTGGATGAGCTGGGACTGAGCCTCACAGATGAGCTGTCAAACCTTCCCTCCACTGGAGG
 CTCACTCAGTGTGGCAGCTGGTGGGAAGAAAGCAGAGGCCACAGCCTCAGCCCTGGCTGATGCTGATGCA
 GACCTGGAAGAGAGGCTCAAGAACCTTCGAGGGACTGA

ACGCGTACGCGGCCGCTCGAGCAGAACTCATCTCAGAAGAGGATCTGGCAGCAAATGATATCCTGGATT
 ACAAGGATGACGACGATAAGGTTTAA

| | |
|--------------------|-----------|
| Restriction Sites: | SgfI-MluI |
| ACCN: | NM_026885 |
| Insert Size: | 669 bp |


[View online »](#)

| | |
|-------------------------------|---------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| OTI Disclaimer: | Our molecular clone sequence data has been matched to the reference identifier above as a point of reference. Note that the complete sequence of our molecular clones may differ from the sequence published for this corresponding reference, e.g., by representing an alternative RNA splicing form or single nucleotide polymorphism (SNP). |
| Components: | The ORF clone is ion-exchange column purified and shipped in a 2D barcoded Matrix tube containing 10ug of transfection-ready, dried plasmid DNA (reconstitute with 100 ul of water). |
| Reconstitution Method: | <ol style="list-style-type: none"> 1. Centrifuge at 5,000xg for 5min. 2. Carefully open the tube and add 100ul of sterile water to dissolve the DNA. 3. Close the tube and incubate for 10 minutes at room temperature. 4. Briefly vortex the tube and then do a quick spin (less than 5000xg) to concentrate the liquid at the bottom. 5. Store the suspended plasmid at -20°C. The DNA is stable for at least one year from date of shipping when stored at -20°C. |
| Note: | Plasmids are not sterile. For experiments where strict sterility is required, filtration with 0.22um filter is required. |
| RefSeq: | NM_026885.3 , NP_081161.1 |
| RefSeq Size: | 927 bp |
| RefSeq ORF: | 669 bp |
| Locus ID: | 68953 |
| UniProt ID: | Q9DB34 |
| Cytogenetics: | 7 |
| Gene Summary: | <p>Probable core component of the endosomal sorting required for transport complex III (ESCRT-III) which is involved in multivesicular bodies (MVBs) formation and sorting of endosomal cargo proteins into MVBs. MVBs contain intraluminal vesicles (ILVs) that are generated by invagination and scission from the limiting membrane of the endosome and mostly are delivered to lysosomes enabling degradation of membrane proteins, such as stimulated growth factor receptors, lysosomal enzymes and lipids. The MVB pathway appears to require the sequential function of ESCRT-O, -I, -II and -III complexes. ESCRT-III proteins mostly dissociate from the invaginating membrane before the ILV is released. The ESCRT machinery also functions in topologically equivalent membrane fission events, such as the terminal stages of cytokinesis. Together with SPAST, the ESCRT-III complex promotes nuclear envelope sealing and mitotic spindle disassembly during late anaphase. ESCRT-III proteins are believed to mediate the necessary vesicle extrusion and/or membrane fission activities, possibly in conjunction with the AAA ATPase VPS4.[UniProtKB/Swiss-Prot Function]</p> <p>Transcript Variant: This variant (2) differs in the 5' UTR compared to variant 1. Variants 1 and 2 both encode the same isoform (a).</p> |