

## Product datasheet for **MC207688**

### **Pcgf1 (NM\_197992) Mouse Untagged Clone**

#### Product data:

**Product Type:** Expression Plasmids  
**Product Name:** Pcgf1 (NM\_197992) Mouse Untagged Clone  
**Tag:** Tag Free  
**Symbol:** Pcgf1  
**Synonyms:** 2010002K04Rik; AU024121; Nspc1  
**Vector:** pCMV6-Entry (PS100001)  
**E. coli Selection:** Kanamycin (25 ug/mL)  
**Cell Selection:** Neomycin  
**Fully Sequenced ORF:** >MC207688 representing NM\_197992  
**Red**=Cloning site **Blue**=ORF **Orange**=Stop codon

TTTTGTAATACGACTCACTATAGGGCGCCGGGAATTCGTCGACTGGATCCGGTACCGAGGAGATCTGCC  
GCC**GCGATCGCC**

ATGAGGCTTCGGAACCAGCTCCAGTCAGTGTACAAGATGGACCCACTACGGAACGAGGAGGAGTCCGGG  
TGAAGATCAAAGACCTGAATGAACACATCGTTTGTCTGCCTGTGCGCGGGCTACTTCGTGGATGCCACCAC  
CATCACAGAGTGTCTCCACACCTTCTGCAAAAGTTGTATCGTGAAGTACCTGCAAACCAGCAAGTACTGC  
CCCATGTGCAACATCAAGATCCACGAGACGCAGCCGTTGCTCAATCTCAAACCTGGATCGGGTCATGCAGG  
ACATAGTGTATAAGCTAGTGCCAGGCTTGAAGACAGTGAAGAGAAACGGATTCCGGGAATTCTATCAGTC  
CCGAGGCTTAGACAGAGTCTCCAGCCAGTGGTGAAGAGCCAGCCCTGAGCAACCTTGGCTCCCTTTC  
AGCAGTTTTGACCACTCTAAAGCCCACTATTATCGATATGATGAACAGCTGAGCCTATGCCTGGAGCGGC  
TGAGTTCTGGCAAAGACAAGAATAAAAAATGCTTTCAGAACAAAGTATGTTTCGGTGTCTGTGAGAGCTGA  
GGTCCGCCATCTCCGAAGGGTCTGTGTCACCGACTAATGCTAAATCCACAGCATGTACAGCTCCTTTTT  
GACAATGAGGTTCTCCAGATCACATGACAATGAAACAGCTATGGCTGTCCCGCTGGTTCGGCAAGCCAT  
CTCCTTTGCTTCTCCAATACAGTGTGAAAGAGAAGAGGAGG**TAG**

**ACGCGT**ACGCGGCCGCTCGAGCAGAAAACCTCATCTCAGAAGAGGATCTGGCAGCAAATGATATCCTGGATT  
ACAAGGATGACGACGATAAGGTTTAA

**Restriction Sites:** SgfI-MluI  
**ACCN:** NM\_197992  
**Insert Size:** 744 bp



[View online »](#)

|                               |  |
|-------------------------------|--|
| <b>OTI Disclaimer:</b>        | Our molecular clone sequence data has been matched to the reference identifier above as a point of reference. Note that the complete sequence of our molecular clones may differ from the sequence published for this corresponding reference, e.g., by representing an alternative RNA splicing form or single nucleotide polymorphism (SNP).   |
| <b>Components:</b>            | The ORF clone is ion-exchange column purified and shipped in a 2D barcoded Matrix tube containing 10ug of transfection-ready, dried plasmid DNA (reconstitute with 100 ul of water).   |
| <b>Reconstitution Method:</b> | <ol style="list-style-type: none"><li>1. Centrifuge at 5,000xg for 5min.</li><li>2. Carefully open the tube and add 100ul of sterile water to dissolve the DNA.</li><li>3. Close the tube and incubate for 10 minutes at room temperature.</li><li>4. Briefly vortex the tube and then do a quick spin (less than 5000xg) to concentrate the liquid at the bottom.</li><li>5. Store the suspended plasmid at -20°C. The DNA is stable for at least one year from date of shipping when stored at -20°C.</li></ol>  |
| <b>RefSeq:</b>                | <a href="#">NM_197992.1</a> , <a href="#">NP_932109.1</a>  |
| <b>RefSeq Size:</b>           | 908 bp   |
| <b>RefSeq ORF:</b>            | 744 bp   |
| <b>Locus ID:</b>              | 69837  |
| <b>UniProt ID:</b>            | <a href="#">Q8R023</a>   |
| <b>Cytogenetics:</b>          | 6 35.94 cM   |
| <b>Gene Summary:</b>          | Component of the Polycomb group (PcG) multiprotein BCOR complex, a complex required to maintain the transcriptionally repressive state of some genes, such as BCL6 and the cyclin-dependent kinase inhibitor, CDKN1A. Transcriptional repressor that may be targeted to the DNA by BCL6; this transcription repressor activity may be related to PKC signaling pathway. Represses CDKN1A expression by binding to its promoter, and this repression is dependent on the retinoic acid response element (RARE element). Promotes cell cycle progression and enhances cell proliferation as well. May have a positive role in tumor cell growth by down-regulating CDKN1A. Component of a Polycomb group (PcG) multiprotein PRC1-like complex, a complex class required to maintain the transcriptionally repressive state of many genes, including Hox genes, throughout development. PcG PRC1 complex acts via chromatin remodeling and modification of histones; it mediates monoubiquitination of histone H2A 'Lys-119', rendering chromatin heritably changed in its expressibility. Within the PRC1-like complex, regulates RNF2 ubiquitin ligase activity. Regulates the expression of DPPA4 and NANOG in the NT2 embryonic carcinoma cells.[UniProtKB/Swiss-Prot Function] |