

Product datasheet for **MC207773**

Rps27a (NM_001033865) Mouse Untagged Clone

Product data:

Product Type: Expression Plasmids
Product Name: Rps27a (NM_001033865) Mouse Untagged Clone
Tag: Tag Free
Symbol: Rps27a
Synonyms: 0610006J14Rik; Uba52; Ubb; Ubc
Vector: pCMV6-Entry (PS100001)
E. coli Selection: Kanamycin (25 ug/mL)
Cell Selection: Neomycin
Fully Sequenced ORF: >MC207773 representing NM_001033865
Red=Cloning site **Blue**=ORF **Orange**=Stop codon

TTTTGTAATACGACTCACTATAGGGCGCCGGAATTCGTCGACTGGATCCGGTACCGAGGAGATCTGCC
GCC**GCGATCGCC**

ATGCAGATCTTTGTGAAGACCC**TACGGGGAAAACCATCAGCTCGAGGTTGAACCCTCGGACACTATAG**
AAAATGTAAGGCCAAGATCCAGGATAAGGAAGGAATTCCTCTGATCAGCAGAGGCTGATCTTTGCTGG
TAAGCAGCTGGAAGATGGCCGACTTTGTCTGACTACAACATTCAAAGGAGTCCACCCTTCATCTGGTG
TTGAGACTTCGGGGTGGTGCTAAGAAAAGGAAGAAGTCTTACACCACTCCCAAGAAGAACAAGCATA
AGAGGAAGAAGGTTAAGTTGGCTGTGCTGAAATACTATAAGGTGGATGAAAATGGCAAAATTAGCCGACT
TCGTCGAGAGTGTCTTCTGATGAATGTGGTGCTGGAGTTTTATGGGAAGCCACTTTGACAGGCATTAC
TGTGGCAAGTGTCTGACTTACTGCTTCAACAAACCAGAAGACAAG**TAG**

ACGCGTACGCGGCCGCTCGAGCAGAACTCATCTCAGAAGAGGATCTGGCAGCAATGATATCCTGGATT
ACAAGGATGACGACGATAAGGTTTAA

Restriction Sites: SgfI-MluI
ACCN: NM_001033865
Insert Size: 471 bp
OTI Disclaimer:

Our molecular clone sequence data has been matched to the reference identifier above as a point of reference. Note that the complete sequence of our molecular clones may differ from the sequence published for this corresponding reference, e.g., by representing an alternative RNA splicing form or single nucleotide polymorphism (SNP).



[View online »](#)

Components:	The ORF clone is ion-exchange column purified and shipped in a 2D barcoded Matrix tube containing 10ug of transfection-ready, dried plasmid DNA (reconstitute with 100 ul of water).
Reconstitution Method:	<ol style="list-style-type: none">1. Centrifuge at 5,000xg for 5min.2. Carefully open the tube and add 100ul of sterile water to dissolve the DNA.3. Close the tube and incubate for 10 minutes at room temperature.4. Briefly vortex the tube and then do a quick spin (less than 5000xg) to concentrate the liquid at the bottom.5. Store the suspended plasmid at -20°C. The DNA is stable for at least one year from date of shipping when stored at -20°C.
RefSeq:	<u>NM_001033865.1</u> , <u>NP_001029037.1</u>
RefSeq Size:	821 bp
RefSeq ORF:	471 bp
Locus ID:	78294
UniProt ID:	<u>P62983</u>
Cytogenetics:	11 A3.3
Gene Summary:	<p>Ubiquitin: Exists either covalently attached to another protein, or free (unanchored). When covalently bound, it is conjugated to target proteins via an isopeptide bond either as a monomer (monoubiquitin), a polymer linked via different Lys residues of the ubiquitin (polyubiquitin chains) or a linear polymer linked via the initiator Met of the ubiquitin (linear polyubiquitin chains). Polyubiquitin chains, when attached to a target protein, have different functions depending on the Lys residue of the ubiquitin that is linked: Lys-6-linked may be involved in DNA repair; Lys-11-linked is involved in ERAD (endoplasmic reticulum-associated degradation) and in cell-cycle regulation; Lys-29-linked is involved in lysosomal degradation; Lys-33-linked is involved in kinase modification; Lys-48-linked is involved in protein degradation via the proteasome; Lys-63-linked is involved in endocytosis, DNA-damage responses as well as in signaling processes leading to activation of the transcription factor NF-kappa-B. Linear polymer chains formed via attachment by the initiator Met lead to cell signaling. Ubiquitin is usually conjugated to Lys residues of target proteins, however, in rare cases, conjugation to Cys or Ser residues has been observed. When polyubiquitin is free (unanchored-polyubiquitin), it also has distinct roles, such as in activation of protein kinases, and in signaling.[UniProtKB/Swiss-Prot Function]</p> <p>Transcript Variant: This variant (2) differs in the 5' UTR, compared to variant 1. Both variants 1 and 2 encode the same protein.</p>