

Product datasheet for MC207793

Eif3m (NM_145380) Mouse Untagged Clone

Product data:

Product Type: Expression Plasmids
Product Name: Eif3m (NM_145380) Mouse Untagged Clone
Tag: Tag Free
Symbol: Eif3m
Synonyms: Ga17; Pcid1; Tango7
Vector: pCMV6-Entry (PS100001)
E. coli Selection: Kanamycin (25 ug/mL)
Cell Selection: Neomycin
Fully Sequenced ORF: >MC207793 representing NM_145380
 Red=Cloning site Blue=ORF Orange=Stop codon

TTTTGTAATACGACTCACTATAGGGCGCCGGAATTCGTCGACTGGATCCGGTACCGAGGAGATCTGCC
 GCC**CGGATCGCC**

ATGAGCGTCCCGCCTTCATCGATATCAGCGAGGAAGACCAGGCTGCTGAACTTCGAGCTTATCTGAAAT
 CAAAAGGAGCTGAGCTCTCAGAAGAAAACCTCGGAAGTGGACTTCATGTAGATTTAGCTCAGATTATTGA
 AGCCTGTGATGTGTGCTGAAGGAAGATGATAAAGATGTTGAAAGTGTGATGAACAGTGTGGTGTCCCTC
 CTGTTGATCCTGGAACCAGACAAGCAAGAAGCATTGATTGAAAGTCTGTGTGAAAAGCTGGTCAAGTTTC
 GGAAGGCGAACCCATCTCTGAGACTACAGTTGCTCAGCAACCTTTTCCATGGGATGGACAAGAATAC
 TCCTGTGAGGTACACAGTGTACTGCAGCCTCATCAAAGTGGCGGCATCATGTGGGGCCATCCAGTACATC
 CCAACGGAGCTGGATCAAGTGAGAAAAGTGGATCTGACTGGAAGCTCACCAGTAAAAGAAGCACACCC
 TCCTCAGACTGCTCTACGAGGCGCTCGTGGATTGTAAGAAAAGTGTGCTGCATCAAAGGTATGGTGGG
 ATTGCTCGGGAGTTACACAGAGGACAATGCTTCCAGGCTCGAGTTGATGCCACAGGTGATTGTGCGG
 GCACTAAAGGATCCAAATGCCTTTCTTTTGACCATCTTCTTACTCTAAAGCCAGTCAAATTTTGAAG
 GCGAGCTTATCCATGATTTATTAACATTTTGTGAGTGCTAAATGGCATCGTATGTCAAGTTCTATCA
 GAATAACAAAGACTTCATAGATTCACCTTGCTTGTACATGAGCAAAATATGGCGAAAATGAGACTCCTC
 ACCTTTATGGAATGGCAGTGGAAAACAAGAGATTTCTTTTGACACAATGCAGCAAGAGCTACAGATTG
 GAGCTGATGATGTTGAAGCATTTGTTATTGATGCTGTAAGAACTAAAATGGTCTACTGCAAAATTGATCA
 GACCCAAAGAAAAGTAGTTGTGAGTACAGCACACATCGGACATTTGAAAACAGCAGTGGCAACAACCTG
 TATGACTCTTAATGCCTGGAAACAGAAGTGAACAAAGTAAAAATAGCCTTTTGAGTCTTTCAGATA
 CCTGA

ACGCGTACGCGGCCGCTCGAGCAGAAAACCTCATCTCAGAAGAGGATCTGGCAGCAAATGATATCCTGGATT
 ACAAGGATGACGACGATAAGGTTTAA

Restriction Sites: SgfI-MluI



[View online »](#)

ACCN:	NM_145380
Insert Size:	1125 bp
OTI Disclaimer:	Our molecular clone sequence data has been matched to the reference identifier above as a point of reference. Note that the complete sequence of our molecular clones may differ from the sequence published for this corresponding reference, e.g., by representing an alternative RNA splicing form or single nucleotide polymorphism (SNP).
Components:	The ORF clone is ion-exchange column purified and shipped in a 2D barcoded Matrix tube containing 10ug of transfection-ready, dried plasmid DNA (reconstitute with 100 ul of water).
Reconstitution Method:	<ol style="list-style-type: none">1. Centrifuge at 5,000xg for 5min.2. Carefully open the tube and add 100ul of sterile water to dissolve the DNA.3. Close the tube and incubate for 10 minutes at room temperature.4. Briefly vortex the tube and then do a quick spin (less than 5000xg) to concentrate the liquid at the bottom.5. Store the suspended plasmid at -20°C. The DNA is stable for at least one year from date of shipping when stored at -20°C.
RefSeq:	<u>NM_145380.2</u> , <u>NP_663355.1</u>
RefSeq Size:	1230 bp
RefSeq ORF:	1125 bp
Locus ID:	98221
UniProt ID:	<u>Q99JX4</u>
Cytogenetics:	2 55.06 cM
Gene Summary:	Component of the eukaryotic translation initiation factor 3 (eIF-3) complex, which is required for several steps in the initiation of protein synthesis. The eIF-3 complex associates with the 40S ribosome and facilitates the recruitment of eIF-1, eIF-1A, eIF-2:GTP:methionyl-tRNA ⁱ and eIF-5 to form the 43S pre-initiation complex (43S PIC). The eIF-3 complex stimulates mRNA recruitment to the 43S PIC and scanning of the mRNA for AUG recognition. The eIF-3 complex is also required for disassembly and recycling of post-termination ribosomal complexes and subsequently prevents premature joining of the 40S and 60S ribosomal subunits prior to initiation. The eIF-3 complex specifically targets and initiates translation of a subset of mRNAs involved in cell proliferation, including cell cycling, differentiation and apoptosis, and uses different modes of RNA stem-loop binding to exert either translational activation or repression.[UniProtKB/Swiss-Prot Function]