

## **Product datasheet for MC208210**

## Calr (NM\_007591) Mouse Untagged Clone

## **Product data:**

**Product Type:** Expression Plasmids

**Product Name:** Calr (NM\_007591) Mouse Untagged Clone

Tag: Tag Free

**Symbol:** Calr

Synonyms: Calregulin; CRT

Mammalian Cell

Selection:

Neomycin

Vector:pCMV6-Entry (PS100001)E. coli Selection:Kanamycin (25 ug/mL)

Restriction Sites: Sgfl-Mlul

**ACCN:** NM\_007591

**Insert Size:** 1251 bp

**OTI Disclaimer:** Our molecular clone sequence data has been matched to the reference identifier above as a

point of reference. Note that the complete sequence of our molecular clones may differ from the sequence published for this corresponding reference, e.g., by representing an alternative

RNA splicing form or single nucleotide polymorphism (SNP).

**Components:** The ORF clone is ion-exchange column purified and shipped in a 2D barcoded Matrix tube

containing 10ug of transfection-ready, dried plasmid DNA (reconstitute with 100 ul of water).

**Reconstitution Method:** 1. Centrifuge at 5,000xg for 5min.

2. Carefully open the tube and add 100ul of sterile water to dissolve the DNA.

3. Close the tube and incubate for 10 minutes at room temperature.

4. Briefly vortex the tube and then do a quick spin (less than 5000xg) to concentrate the liquid

at the bottom.

5. Store the suspended plasmid at -20°C. The DNA is stable for at least one year from date of

shipping when stored at -20°C.

**RefSeq:** NM 007591.3, NP 031617.1

RefSeq Size: 1943 bp RefSeq ORF: 1251 bp



**OriGene Technologies, Inc.** 9620 Medical Center Drive, Ste 200

CN: techsupport@origene.cn

Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com



## Calr (NM\_007591) Mouse Untagged Clone - MC208210

**Locus ID:** 12317

 UniProt ID:
 P14211

 Cytogenetics:
 8 41.21 cM

Gene Summary: Calcium-binding chaperone that promotes folding, oligomeric assembly and quality control in

the endoplasmic reticulum (ER) via the calreticulin/calnexin cycle. This lectin interacts

transiently with almost all of the monoglucosylated glycoproteins that are synthesized in the

ER. Interacts with the DNA-binding domain of NR3C1 and mediates its nuclear export.

Involved in maternal gene expression regulation. May participate in oocyte maturation via the  $\,$ 

regulation of calcium homeostasis (By similarity).[UniProtKB/Swiss-Prot Function]