

Product datasheet for MC208267

Cd8a (NM_009857) Mouse Untagged Clone

Product data:

Product Type:	Expression Plasmids
Product Name:	Cd8a (NM_009857) Mouse Untagged Clone
Tag:	Tag Free
Symbol:	Cd8a
Synonyms:	BB154331; Ly-2; Ly-35; Ly-B; Lyt-2
Mammalian Cell Selection:	Neomycin
Vector:	pCMV6-Entry (PS100001)
E. coli Selection:	Kanamycin (25 ug/mL)
Fully Sequenced ORF:	>MC208267 representing NM_009857 Red=Cloning site Blue=ORF Orange=Stop codon

TTTGTAAATACGACTCACTATAGGGCGGCCGGAATTCGTCGACTGGATCCGGTACCGAGGAGATCTGCC
 GCCCGCATCGCC

ATGGCCTCACCGTTGACCCGCTTTCTGTCGCTGAACCTGCTGCTGCTGGGTGAGTCGATTATCCTGGGGA
 GTGGAGAAGCTAAGCCACAGGCACCCGAATCCGAATCTTCCAAAGAAAATGGACGCCGAACCTGGTCA
 GAAGGTGGACCTGGTATGTGAAGTGTGGGGTCCGTTTCGCAAGGATGCTCTTGGCTCTTCCAGAACTCC
 AGCTCCAAACTCCCCAGCCACCTTCGTTGTCTATATGGCTTCATCCACAACAAGATAACGTGGGACG
 AGAAGCTGAATTCGTCGAACTGTTTTCTGCCATGAGGGACACGAATAATAAGTACGTTCTACCCCTGAA
 CAAGTTCAGCAAGGAAAACGAAGGCTACTATTTCTGCTCAGTCATCAGCAACTCGGTGATGACTTCAGT
 TCTGTCGTGCCAGTCTTCAGAAAGTGAACCTACTACTACCAAGCCAGTGTGCGAACTCCCTCACCTG
 TGCACCCTACCGGGACATCTCAGCCCCAGAGACCAGAAGATTGTCGGCCCCGTGGCTCAGTGAAGGGGAC
 CGGATTGGACTTCGCCTGTGATATTTACATCTGGGCACCCTTGGCCGGAATCTGCGTGGCCCTTCTGCTG
 TCCTTGATCATCACTCTCATCTGCTACCACAGCCGCTAG

ACGCGTACGCGGCCGCTCGAGCAGAAACTCATCTCAGAAGAGGATCTGGCAGCAAATGATATCCTGGATT
 ACAAGGATGACGACGATAAGGTTTAA

Restriction Sites:	SgfI-MluI
ACCN:	NM_009857
Insert Size:	669 bp


[View online »](#)

OTI Disclaimer:	Our molecular clone sequence data has been matched to the reference identifier above as a point of reference. Note that the complete sequence of our molecular clones may differ from the sequence published for this corresponding reference, e.g., by representing an alternative RNA splicing form or single nucleotide polymorphism (SNP).
Components:	The ORF clone is ion-exchange column purified and shipped in a 2D barcoded Matrix tube containing 10ug of transfection-ready, dried plasmid DNA (reconstitute with 100 ul of water).
Reconstitution Method:	<ol style="list-style-type: none"> 1. Centrifuge at 5,000xg for 5min. 2. Carefully open the tube and add 100ul of sterile water to dissolve the DNA. 3. Close the tube and incubate for 10 minutes at room temperature. 4. Briefly vortex the tube and then do a quick spin (less than 5000xg) to concentrate the liquid at the bottom. 5. Store the suspended plasmid at -20°C. The DNA is stable for at least one year from date of shipping when stored at -20°C.
Note:	Plasmids are not sterile. For experiments where strict sterility is required, filtration with 0.22um filter is required.
RefSeq:	NM_009857.1 , NP_033987.1
RefSeq Size:	3099 bp
RefSeq ORF:	669 bp
Locus ID:	12525
Cytogenetics:	6 32.14 cM
Gene Summary:	<p>Integral membrane glycoprotein that plays an essential role in the immune response and serves multiple functions in responses against both external and internal offenses. In T-cells, functions primarily as a coreceptor for MHC class I molecule:peptide complex. The antigens presented by class I peptides are derived from cytosolic proteins while class II derived from extracellular proteins. Interacts simultaneously with the T-cell receptor (TCR) and the MHC class I proteins presented by antigen presenting cells (APCs). In turn, recruits the Src kinase LCK to the vicinity of the TCR-CD3 complex. LCK then initiates different intracellular signaling pathways by phosphorylating various substrates ultimately leading to lymphokine production, motility, adhesion and activation of cytotoxic T-lymphocytes (CTLs). This mechanism enables CTLs to recognize and eliminate infected cells and tumor cells. In NK-cells, the presence of CD8A homodimers at the cell surface provides a survival mechanism allowing conjugation and lysis of multiple target cells. CD8A homodimer molecules also promote the survival and differentiation of activated lymphocytes into memory CD8 T-cells.[UniProtKB/Swiss-Prot Function]</p> <p>Transcript Variant: This variant (2) lacks an exon in the coding region resulting in a frameshift, compared to variant 1. It encodes isoform 2, which is shorter and has a distinct C-terminus, compared to isoform 1. Sequence Note: The RefSeq transcript and protein were derived from genomic sequence to make the sequence consistent with the reference genome assembly. The genomic coordinates used for the transcript record were based on alignments.</p>