

## **Product datasheet for MC208280**

## Cflar (NM 009805) Mouse Untagged Clone

**Product data:** 

**Product Type:** Expression Plasmids

**Product Name:** Cflar (NM\_009805) Mouse Untagged Clone

Tag: Tag Free Symbol: Cflar

Synonyms: 2310024N18Rik; A430105C05Rik; c-Flip; Cash; Casper; CLARP; ENSMUSG00000072980; FLAME;

FLAME-1; Flip

Mammalian Cell

Selection:

Neomycin

Vector:pCMV6-Entry (PS100001)E. coli Selection:Kanamycin (25 ug/mL)

Fully Sequenced ORF: >MC208280 representing NM\_009805

Red=Cloning site Blue=ORF Orange=Stop codon

TTTTGTAATACGACTCACTATAGGGCGGCCGGGAATTCGTCGACTGGATCCGGTACCGAGGAGATCTGCC

GCCGCGATCGCC

ATGGCCCAGAGCCCTGTGTCTGCCGAGGTCATTCACCAGGTGGAAGAGTGTCTTGATGAAGACGAGAAGG
AGATGATGCTCTTCCTGTGTAGAGATGTGACTGAGAACCTGGCTGCACCTAACGTCAGGGACCTCCTGGA
TAGCTTAAGTGAGAGAGGCCAGCTCTCTTTTGCTACCTTGGCTGAATTGCTCTACAGAGTGAGGCGGTTT
GACCTTCTCAAGAGGATCTTGAAGACAGACAAAGCAACCGTGGAGGACCACCTGCGCAGAAACCCTCACC
TGGTTTCTGATTATAGGGTCCTGCTGATGGAGGATTGGTGAGAGCCTTAGATCAGAACGATGTATCCTCCTT
AGTTTTCCTTACAAGGGATTACACAGGCAGAGGCAAGATAGCCAAGGACAAGAGTTTCTTGGATCTGGTG
ATTGAATTGGAGAAACTGAATCTAATTGCTTCAGACCAATTGAATTTGTTAGAAAAAATGCCTGAAGAACA
TCCACAGAATAGACTTGAACACAAAGATCCAGAAGTACACCCAGTCCAGCCAAGGAGCAAGATCAAATAT
GAATACTCTCCAGGCTTCGCTCCCAAAAATTGAGTATCAAGTATAACTCAAGGGTGAGTCTGGAGCCAGTG
TATGGAGTACCAGCATGA

**ACGCGT**ACGCGGCCGCTCGAGCAGAAACTCATCTCAGAAGAGGATCTGGCAGCAAATGATATCCTGGATT

ACAAGGATGACGACGATAAGGTTTAA

Chromatograms: <a href="https://cdn.origene.com/chromatograms/ja1875">https://cdn.origene.com/chromatograms/ja1875</a> b11.zip

**Restriction Sites:** Sgfl-Mlul ACCN: NM\_009805

**Insert Size:** 648 bp



**OriGene Technologies, Inc.** 9620 Medical Center Drive, Ste 200

CN: techsupport@origene.cn

Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com



## **OTI Disclaimer:**

Due to the inherent nature of this plasmid, standard methods to replicate additional amounts of DNA in E. coli are highly likely to result in mutations and/or rearrangements. Therefore, OriGene does not guarantee the capability to replicate this plasmid DNA. Additional amounts of DNA can be purchased from OriGene with batch-specific, full-sequence verification at a reduced cost. Please contact our customer care team at <a href="mailto:customercom">customercom</a> or by calling 301.340.3188 option 3 for pricing and delivery.

The molecular sequence of this clone aligns with the gene accession number as a point of reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing variants is recommended prior to use. <u>More info</u>

Components:

The ORF clone is ion-exchange column purified and shipped in a 2D barcoded Matrix tube containing 10ug of transfection-ready, dried plasmid DNA (reconstitute with 100 ul of water).

**Reconstitution Method:** 

- 1. Centrifuge at 5,000xg for 5min.
- 2. Carefully open the tube and add 100ul of sterile water to dissolve the DNA.
- 3. Close the tube and incubate for 10 minutes at room temperature.
- 4. Briefly vortex the tube and then do a quick spin (less than 5000xg) to concentrate the liquid at the bottom.
- 5. Store the suspended plasmid at -20°C. The DNA is stable for at least one year from date of shipping when stored at -20°C.

Note:

Plasmids are not sterile. For experiments where strict sterility is required, filtration with 0.22um filter is required.

**RefSeq:** NM 009805.4, NP 033935.2

RefSeq Size:2036 bpRefSeq ORF:648 bpLocus ID:12633

Cytogenetics: 1 29.16 cM

**Gene Summary:** 

Apoptosis regulator protein which may function as a crucial link between cell survival and cell death pathways in mammalian cells. Acts as an inhibitor of TNFRSF6 mediated apoptosis. A proteolytic fragment (p43) is likely retained in the death-inducing signaling complex (DISC) thereby blocking further recruitment and processing of caspase-8 at the complex. Full length and shorter isoforms have been shown either to induce apoptosis or to reduce TNFRSF-triggered apoptosis. Lacks enzymatic (caspase) activity (By similarity).[UniProtKB/Swiss-ProtFunction]