

Product datasheet for **MC208391**

Dio2 (NM_010050) Mouse Untagged Clone

Product data:

Product Type:	Expression Plasmids
Product Name:	Dio2 (NM_010050) Mouse Untagged Clone
Symbol:	Dio2
Synonyms:	5DII; AI324267; DIOII
Mammalian Cell Selection:	Neomycin
Vector:	pCMV6-Entry (PS100001)
E. coli Selection:	Kanamycin (25 ug/mL)



[View online »](#)

Fully Sequenced ORF: >NCBI ORF sequence for NM_010050, the custom clone sequence may differ by one or more nucleotides

```

TGACTGGGGAAGCAGAGTGCCAGGAGACTGACCGAGGAGGCAGAGAAGATGGGACTCCTCAGCGTAGAC
TTGCTGATCACCTCGCATCCTGCCAGTCTTTTTCTCCAACGCCTCTCTGGCGCTCTATGACTCGG
TCATTCTGCTCAAGCACGTGGCGTTGCTTCTGAGCCGCTCCAAGTCCACTCGCGGAGAGTGGAGGCGCAT
GCTGACCTCAGAAGGGCTGCGCTGTGTCTGGAACAGCTTCTCCTAGATGCCTACAAACAGGTTAAACTG
GGTGAAGATGCTCCCAATTCCAGTGTGGTGCACGTCTCCAATCCTGAATCAGGTAACAATTATGCCTCGG
AGAAGACCGCTGATGGGGCCGAATGCCACCTTCTTGACTTTGCCAGTGCAGAGCGCCCACTGGTGGTCAA
CTTTGGTTCAGCCACCTGACCACCTTCTACTAGGCAACTGCCAGCCTTCCGCCAGCTGGTGAAGAGTTC
TCCTCGTGGTACTTCTGTTGGTATACATTGATGAGGCTCACCTTCGGATGGCTGGCAGTGCCTG
GGGACTCCTCTGTCTTTGAGGTTAAGAAGCACCGGAACCAAGAGGACCGATGTGCAGCAGCTCACCA
GCTCCTGGAGCGTTTCTCCTTGGCCGCCAGTGTCAAGTTGTGGTGCACCGCATGGACAATAATGCCAAC
GTAGCTTACGGGGTAGCCTTTGACGTGTGTGCATCGTGCAGAGACGGAAAATTGCTTACTTAGGAGGGA
AGGGCCCTTTCAGCTATAACCTACAAGAAGTCCGAAGTTGGCTGGAGAAGAAATTCAGCAAGAGATGAAT
TCTAGATTAGCTGGATACAAGTGTGATTGTAATAGAGTTTATTATTTAAAGAAATATGTAAAAGAAGT
GAGAACTGAACTGAATCTATTATTTCAACTGAATCCCATTGCCTCACGAAAGACGGGGACACACCTGTC
AGAAGACCTTGAACCTCTTAACATCTCAATACTTTCTCTAGTACACAAATGACATCTGGCTAAATAGCAG
CCCTTTTACCTCCCTTCTGAGCGAATTGATCCAAACTGGAGTGCAAGAAACCCTGTTGCAGCCTGTATTC
ATTCTTGCTTGCAGAAAGAAGTATTCTGTCTGCGTGTGCTCTGGGACCAGAGGACCATAAGTTCTCAGA
ATACTTATTGTTAAGGGAGCTACTACCTATGATCTGATTAAGTGTATTGTCTCAGCTTTAGTCACCAGAA
ACAAACCAACTGGAAAGTCCCCAGATACATACTACGATGTGATCTTGAAGCCATGCCATGAGTTAG
GACTGAAGTGTTAGTTCAGCCATGCAATGTCCGGTTAGTCTCTGTAGCCCCGAATTCCTCAAGTGTGAC
AAGCAAAGACAGAGGAAGCAGAAAGATGAGGAAATCGCATAGTGTTTTTACCTGGAAGCTGTAGGTAATG
AGTGTGTTCAACAGCATCAGCATTGAGGCATGTACACAGCTGGGAAAACAAAGGCGATAACTAAAAGTA
GCACGGGCAGGATGCACACGAAAGGAGATGTGTAGCAATATTCCTACACTGAAAGGAAGTGGGCGATGG
CAAAGATAGGTGAACGTTTTAATGTGGAGAGAAGTACTTTATGGGACTAATTGAGATATACCCATTTCA
GTGGATAATATCCGTTCCAGAGAAAACCTCCAGGGGTCTGGTGTGATGTGTTGTTCTGCCAAATCCTA
CAGTCACAATGCTACACCAGCTCCATTCTAACCTCTGTAGCATTGTTTTCTTTGGGTGCTGAGTGAAG
GGTGTAAAGAGGCAGGTACACTAACCATTAGTTTCTGCCTTCTCCTCCAGTAAAAGAAGTGAATGATA
AGGGATTTCTTACATAAACATGAGAGCAATTTATTCTGAGAATCTCTACCTATAGTTTTGAGGGCTTAC
ACAAGCTTATTCTTGGTACCACAGGAATTTGGTTAAATGGAATTTCTTTGACATTTGAATTTGACTTG
GTTCTTTTTTTGTTCAACTGGGATTAACCCATTAGTTGACACTTAATAGCTGTGTACATACCTATGATAAA
AACTAGATATGAATACATATCTCTTAAAAATAACAAGAGACACAGCTGTACCTACAAGCAATACCTAAAGAT
CACAAATAAAAAACACTCCACAAAAAACCCAAAAACCAACCAACAGATCAACCAATGATTGTA
ATTATTTTTTAACTCAAAGACATTAAGATCTCAACATATGGTTATAGTTACATGCCAAATCAAGGGTCT
TATGCTATTCCATTCAATAATTAACCATATATTGAGTTATTTGAAAAAGACAATTGAAATTTAAAA
GTACATTTGTTAAAAATATATTGTGCAAAATGTGGAAAATAGGAAGTGGGTATGCATAAAACAAGAGTCTTG
TTTTTAAAAATGGGAGCAAGAAAATAAGTCAGATCGCTAACCCATATATTGTTTATGTCATGGAAGCAGCAC
TATGGCTAATGTTTTTGGCAGCATCTAAGTTCTCCAGTGTTTTAAATTCTGTAAGACACACCCACAAG
GAAACACAGTGAATCCCAAAATGCAACAGTGTCTAATGAACTTCTGTAGAATCATATTGAAGATCGATT
ACTTCAGACTCTCAGATGTCAAGGCCTTCACTTGAGGCTTATGAGCGATCCCTCACTCATCTTCATCTT
CAAAATGCTGTTGCGATTGATGTGGCTCCCTAAAGAAAGTAAAATGGAAAATTTGGCTGCCCCACCCACA
CCGTCGTCGCAAAATGACCCCTTTGGTTCCACGTGTGATAGCGAAGCAGAAAGTCAAGAGTGGGTGGG
GGGATGAGTTTTGGCTAACGCCATGCTAGTTGTCTGGG
    
```

Restriction Sites: Sgfl-MluI
ACCN: NM_010050

OTI Disclaimer:	Our molecular clone sequence data has been matched to the reference identifier above as a point of reference. Note that the complete sequence of our molecular clones may differ from the sequence published for this corresponding reference, e.g., by representing an alternative RNA splicing form or single nucleotide polymorphism (SNP). The expression of this clone is not guaranteed due to the nature of selenoproteins.
OTI Annotation:	This clone encodes a selenoprotein containing the rare amino acid selenocysteine (Sec). Sec is encoded by UGA codon, which normally signals translational termination. Expression of this clone is not guaranteed due to the nature of selenoproteins.
Components:	The ORF clone is ion-exchange column purified and shipped in a 2D barcoded Matrix tube containing 10ug of transfection-ready, dried plasmid DNA (reconstitute with 100 ul of water).
Reconstitution Method:	<ol style="list-style-type: none">1. Centrifuge at 5,000xg for 5min.2. Carefully open the tube and add 100ul of sterile water to dissolve the DNA.3. Close the tube and incubate for 10 minutes at room temperature.4. Briefly vortex the tube and then do a quick spin (less than 5000xg) to concentrate the liquid at the bottom.5. Store the suspended plasmid at -20°C. The DNA is stable for at least one year from date of shipping when stored at -20°C.
RefSeq:	BC125383 , AAI25384
RefSeq Size:	2908 bp
RefSeq ORF:	789 bp
Locus ID:	13371
UniProt ID:	Q9Z1Y9
Cytogenetics:	12 D3

Gene Summary:

The protein encoded by this gene belongs to the iodothyronine deiodinase family. It catalyzes the conversion of prohormone thyroxine (3,5,3',5'-tetraiodothyronine, T4) to the bioactive thyroid hormone (3,5,3'-triiodothyronine, T3) by outer ring 5'-deiodination. This gene is highly expressed in brain, placenta and mammary gland. It is thought to be responsible for the 'local' production of T3, and thus important in influencing thyroid hormone action in these tissues. Knockout studies in mice suggest that this gene may play an important role in brown adipose tissue lipogenesis, auditory function, and bone formation. This protein is a selenoprotein containing the non-standard amino acid, selenocysteine (Sec), which is encoded by the UGA codon that normally signals translation termination. The 3' UTRs of selenoprotein mRNAs contain a conserved stem-loop structure, designated the Sec insertion sequence (SECIS) element, that is necessary for the recognition of UGA as a Sec codon, rather than as a stop signal. Unlike the other two members (DIO1 and DIO3) of this enzyme family, the mRNA for this gene contains an additional in-frame UGA codon that has been reported (in human) to function either as a Sec or a stop codon, resulting in two potential isoforms with one or two Sec residues; however, only the upstream Sec (conserved with the single Sec residue found at the active site in DIO1 and DIO3) was shown to be essential for enzyme activity (PMID:10403186). In addition, the lack of conservation of the protein extension past the second TGA codon suggests that the one-Sec containing isoform represents the canonical form. [provided by RefSeq, Oct 2018]