

Product datasheet for **MC208473**

Fcgr2b (NM_001077189) Mouse Untagged Clone

Product data:

Product Type:	Expression Plasmids
Product Name:	Fcgr2b (NM_001077189) Mouse Untagged Clone
Tag:	Tag Free
Symbol:	Fcgr2b
Synonyms:	A1528646; CD32; F630109E10Rik; Fcgr2; Fcgr2a; FcgRII; Fcr-2; Fcr-3; fcRII; Fc[g]RII; Ly-17
Mammalian Cell Selection:	Neomycin
Vector:	pCMV6-Entry (PS100001)
E. coli Selection:	Kanamycin (25 ug/mL)
Fully Sequenced ORF:	>MC208473 representing NM_001077189 Red=Cloning site Blue=ORF Orange=Stop codon

TTTTGTAATACGACTCACTATAGGGCGCCGGGAATTCGTCGACTGGATCCGGTACCGAGGAGATCTGCC
GCC**CGATCGCC**

ATGGGAATCCTGCCGTTCTACTGATCCCCATGGAGAGCAACTGGACTGTCCATGTGTTCTCACGGACTT
TGTGCCATATGCTACTGTGGACAGCCGTGCTAAATCTTGCTGCTGGGACTCATGATCTTCCAAAGGCTGT
GGTCAAACCTCGAGCCCCGTGGATCCAGGTGCTCAAGGAAGACACGGTGACACTGACATGCGAAGGGACC
CACAACCCTGGAACTCTTCTACCCAGTGGTCCACAATGGGAGGTCCATCCGGAGCCAGGTCCAAGCCA
GCTACACGTTTAAAGGCCACAGTCAATGACAGTGGAGAATATCGGTGTCAAATGGAGCAGACCCGCTCAG
CGACCCTGTAGATCTGGGAGTGATTTCTGACTGGCTGCTGCTCCAGACCCTCAGCTGGTGTCTTCTGGAA
GGGGAAACCATCACGCTAAGGTGCCATAGCTGGAGGAACAACTACTGAACAGGATCTCGTTCTCCATA
ATGAAAAATCCGTGAGGTATCATCACTACAGTAGTAATTTCTCTATCCAAAAGCCAACCACAGTCACAG
TGGGGACTACTACTGCAAAGGAAGTCTAGGAAGGACACTGCACCAGTCCAAGCCTGTCCACACTACTGTC
CAAGGGCCCAAGTCCAGCAGGTCTTTACCAGTATTGACAATTGTGGCTGCTGCTACTGGGATTGCTGTCG
CAGCCATTGTTATTATCCTAGTATCCTTGGTCTATCTCAAGAAAAAGCAGGTTCCAGCTCTCCAGGAAA
CCCTGATCACAGGAAAATGGGAGAAACCCTTCCAGAGGAAGTAGGTGAGTACAGACAGCCCTCTGGGGGC
TCAGTGCTGTACAGCCAGGGCCTCCATCTGGACTGGAGCAACAAGCAGCAGCCCATACAATCCTCCTG
ATCTGGAAGAAGCTGCCAAAAGTGGGCTGAGAATACGATCACCTACTCACTTCTCAAGCATCCCGAAGC
CCTGGATGAAGAAACAGAGCATGATTACCAGAACCACATT**TAG**

ACGCGTACGCGGCCGCTCGAGCAGAAAACCTCATCTCAGAAGAGGATCTGGCAGCAAATGATATCCTGGATT
ACAAGGATGACGACGATAAGGTTTAA

Chromatograms: https://cdn.origene.com/chromatograms/ja2551_d01.zip



[View online »](#)

Restriction Sites:	Sgfl-Mlul
ACCN:	NM_001077189
Insert Size:	1023 bp
OTI Disclaimer:	<p>Due to the inherent nature of this plasmid, standard methods to replicate additional amounts of DNA in E. coli are highly likely to result in mutations and/or rearrangements. Therefore, OriGene does not guarantee the capability to replicate this plasmid DNA. Additional amounts of DNA can be purchased from OriGene with batch-specific, full-sequence verification at a reduced cost. Please contact our customer care team at custsupport@origene.com or by calling 301.340.3188 option 3 for pricing and delivery.</p> <p>The molecular sequence of this clone aligns with the gene accession number as a point of reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing variants is recommended prior to use. More info</p>
Components:	The ORF clone is ion-exchange column purified and shipped in a 2D barcoded Matrix tube containing 10ug of transfection-ready, dried plasmid DNA (reconstitute with 100 ul of water).
Reconstitution Method:	<ol style="list-style-type: none"> 1. Centrifuge at 5,000xg for 5min. 2. Carefully open the tube and add 100ul of sterile water to dissolve the DNA. 3. Close the tube and incubate for 10 minutes at room temperature. 4. Briefly vortex the tube and then do a quick spin (less than 5000xg) to concentrate the liquid at the bottom. 5. Store the suspended plasmid at -20°C. The DNA is stable for at least one year from date of shipping when stored at -20°C.
RefSeq:	NM_001077189.1 , NP_001070657.1
RefSeq Size:	1556 bp
RefSeq ORF:	1023 bp
Locus ID:	14130
Cytogenetics:	1 78.02 cM
Gene Summary:	<p>Receptor for the Fc region of complexed immunoglobulins gamma. Low affinity receptor. Involved in a variety of effector and regulatory functions such as phagocytosis of antigen-antibody complexes from the circulation and modulation of antibody production by B-cells. Isoform IIB1 and isoform IIB1' form caps but fail to mediate endocytosis or phagocytosis. Isoform IIB2 can mediate the endocytosis of soluble immune complexes via clathrin-coated pits. Isoform IIB1 and isoform IIB2 can down-regulate B-cell, T-cell, and mast cell activation when coaggregated to B-cell receptors for AG (BCR), T-cell receptors for AG (TCR), and Fc receptors, respectively.[UniProtKB/Swiss-Prot Function]</p>