

Product datasheet for **MC208772**

II15ra (NM_133836) Mouse Untagged Clone

Product data:

Product Type:	Expression Plasmids
Product Name:	II15ra (NM_133836) Mouse Untagged Clone
Tag:	Tag Free
Symbol:	II15ra
Synonyms:	AA690181; IL-15RA
Mammalian Cell Selection:	Neomycin
Vector:	pCMV6-Entry (PS100001)
E. coli Selection:	Kanamycin (25 ug/mL)
Restriction Sites:	Sgfl-Mlul
ACCN:	NM_133836
Insert Size:	372 bp
OTI Disclaimer:	Our molecular clone sequence data has been matched to the reference identifier above as a point of reference. Note that the complete sequence of our molecular clones may differ from the sequence published for this corresponding reference, e.g., by representing an alternative RNA splicing form or single nucleotide polymorphism (SNP).
Components:	The ORF clone is ion-exchange column purified and shipped in a 2D barcoded Matrix tube containing 10ug of transfection-ready, dried plasmid DNA (reconstitute with 100 ul of water).
Reconstitution Method:	<ol style="list-style-type: none">1. Centrifuge at 5,000xg for 5min.2. Carefully open the tube and add 100ul of sterile water to dissolve the DNA.3. Close the tube and incubate for 10 minutes at room temperature.4. Briefly vortex the tube and then do a quick spin (less than 5000xg) to concentrate the liquid at the bottom.5. Store the suspended plasmid at -20°C. The DNA is stable for at least one year from date of shipping when stored at -20°C.
RefSeq:	<u>NM_133836.2</u> , <u>NP_598597.1</u>
RefSeq Size:	1717 bp
RefSeq ORF:	372 bp



[View online »](#)

Locus ID: 16169

UniProt ID: [Q60819](#)

Cytogenetics: 2 8.97 cM

Gene Summary: High-affinity receptor for interleukin-15 (PubMed:17947230). Can signal both in cis and trans where IL15R from one subset of cells presents IL15 to neighboring IL2RG-expressing cells (PubMed:17947230). In neutrophils, binds and activates kinase SYK in response to IL15 stimulation (By similarity). In neutrophils, required for IL15-induced phagocytosis in a SYK-dependent manner (By similarity).[UniProtKB/Swiss-Prot Function]
Transcript Variant: This variant (2) contains an alternate 5'-most exon in the 5' UTR compared to variant 1. This difference causes translation initiation at a downstream AUG and results in an isoform (2) with a shorter N-terminus, compared to isoform 1. Both variants 2 and 3 encode the same isoform (2). Sequence Note: The RefSeq transcript and protein were derived from genomic sequence to make the sequence consistent with the reference genome assembly. The genomic coordinates used for the transcript record were based on alignments.