

Product datasheet for MC208863

Lamp2 (NM_001017959) Mouse Untagged Clone

Product data:

Product Type:	Expression Plasmids
Product Name:	Lamp2 (NM_001017959) Mouse Untagged Clone
Tag:	Tag Free
Symbol:	Lamp2
Synonyms:	CD107b; Lamp-2; Lamp-2a; Lamp-2b; Lamp-2c; Lamp II; LGP-B; Mac3
Mammalian Cell Selection:	Neomycin
Vector:	pCMV6-Entry (PS100001)
E. coli Selection:	Kanamycin (25 ug/mL)
Fully Sequenced ORF:	>MC208863 representing NM_001017959 Red=Cloning site Blue=ORF Green=Tags(s)

TTTTGTAATACGACTCACTATAGGGCGGCCGGAATTCGTCGACTGGATCCGGTACCGAGGAGATCTGCC
 GCCGCGATCGCC

ATGTGCCTCTCTCCGTTAAAGGCGCAAAGCTCATCTGATCTTTCTGTTCTAGGAGCCGTTTCAGTCCA
 ATGCATTGATAGTTAATTTGACAGATTCAAAGGGTACTTGCTTTATGCAGAATGGGAGATGAATTTAC
 AATAACATATGAACTACAAACCAACCAATAAACTATAACCATTGCAGTACCTGACAAGGCGACACAC
 GATGGAAGCAGTTGTGGGATGACCGGAATAGTGCCAAAATAATGATACAATTTGGATTCTGTCTCTT
 GGGCTGTGAATTTACCAAGGAAGCATCTCATTATTCAATTCATGACATCGTCTTCTACAACACTAG
 TGATAGCACAGTATTTCTGCTGTAGCTAAAGGAGTTCATACTGTTAAAAATCCTGAGAATTTCAAA
 GTTCCATTGGATGTCATCTTTAAGTGCAATAGTGTTTAACTTACAACCTGACTCCTGCTCGTTTCAGAAAT
 ATTGGGGTATTACCTGCAAGCTTTTGCCAAAATGGTACAGTGAGTAAAAATGAACAAGTGTGTGAAGA
 AGACCAAACCTCCCACTGTGGCACCACATTCACACCACTGCCCCGTCGACTACAACCTACACTCACT
 CCAACTTCAACACCACTCCAACCTCCAACCTCCAACCGTTGGAACCTACAGCATTAGAAATGGCA
 ATACTACCTGTCTGCTGGCTACCATGGGCTGCAGCTGAACATCACTGAGGAGAAGGTGCCTTTTCATTTT
 TAACATCAACCCCTGCCACAACCACTTCACCGGCAGCTGTCAACCTCAAAGTGCTCAACTTAGGCTGAAC
 AACAGCCAAATTAAGTATCTTGACTTTATCTTTGCTGTGAAAAATGAAAAACGGTTCTATCTGAAGGAAG
 TGAATGTCTACATGTATTTGGCTAATGGCTCAGCTTTCAACATTTCCAACAAGAACCTTAGCTTCTGGGA
 TGCCCCCTCGGGAAGTTCTTATATGTGCAACAAAGAGCAGGTGCTTTCTGTGCTAGAGCGTTTCAGATC
 AACACCTTTAACCTAAAGGTGCAACCTTTTAAATGTGACAAAAGGACAGTATTCTACAGCTCAAGACTGCA
 GTGCAGATGAAGACAACCTTCCTTGTGCCATAGCGGTGGGAGCAGCTCTGGGAGGAGTACTTATTCTAGT
 GTTGCTGGCTATTTTATTGGTCTCAAGCGCCATCATACTGGATATGAGCAATTTAG

ACGCGTACGCGGCCGCTCGAGCAGAACTCATCTCAGAAGAGGATCTGGCAGCAATGATATCTGGATT
 ACAAGGATGACGACGATAAGGTTTAA


[View online »](#)

Chromatograms:	https://cdn.origene.com/chromatograms/ja1232_d12.zip
Restriction Sites:	Sgfl-MluI
ACCN:	NM_001017959
Insert Size:	1248 bp
OTI Disclaimer:	<p>Due to the inherent nature of this plasmid, standard methods to replicate additional amounts of DNA in E. coli are highly likely to result in mutations and/or rearrangements. Therefore, OriGene does not guarantee the capability to replicate this plasmid DNA. Additional amounts of DNA can be purchased from OriGene with batch-specific, full-sequence verification at a reduced cost. Please contact our customer care team at custsupport@origene.com or by calling 301.340.3188 option 3 for pricing and delivery.</p> <p>The molecular sequence of this clone aligns with the gene accession number as a point of reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing variants is recommended prior to use. More info</p>
Components:	The ORF clone is ion-exchange column purified and shipped in a 2D barcoded Matrix tube containing 10ug of transfection-ready, dried plasmid DNA (reconstitute with 100 ul of water).
Reconstitution Method:	<ol style="list-style-type: none"> 1. Centrifuge at 5,000xg for 5min. 2. Carefully open the tube and add 100ul of sterile water to dissolve the DNA. 3. Close the tube and incubate for 10 minutes at room temperature. 4. Briefly vortex the tube and then do a quick spin (less than 5000xg) to concentrate the liquid at the bottom. 5. Store the suspended plasmid at -20°C. The DNA is stable for at least one year from date of shipping when stored at -20°C.
Note:	Plasmids are not sterile. For experiments where strict sterility is required, filtration with 0.22um filter is required.
RefSeq:	NM_001017959.1 , NP_001017959.1
RefSeq Size:	1768 bp
RefSeq ORF:	1248 bp
Locus ID:	16784
UniProt ID:	P17047
Cytogenetics:	X 22.67 cM

Gene Summary:

Plays an important role in chaperone-mediated autophagy, a process that mediates lysosomal degradation of proteins in response to various stresses and as part of the normal turnover of proteins with a long biological half-life (PubMed:10972293). Functions by binding target proteins, such as GAPDH and MLLT11, and targeting them for lysosomal degradation (By similarity). Required for the fusion of autophagosomes with lysosomes during autophagy (PubMed:27628032). Cells that lack LAMP2 express normal levels of VAMP8, but fail to accumulate STX17 on autophagosomes, which is the most likely explanation for the lack of fusion between autophagosomes and lysosomes (PubMed:27628032). Required for normal degradation of the contents of autophagosomes (PubMed:10972293, PubMed:12221139). Plays a role in lysosomal protein degradation in response to starvation (PubMed:27628032). Required for efficient MHCII-mediated presentation of exogenous antigens via its function in lysosomal protein degradation; antigenic peptides generated by proteases in the endosomal/lysosomal compartment are captured by nascent MHCII subunits. Is not required for efficient MHCII-mediated presentation of endogenous antigens (By similarity). [UniProtKB/Swiss-Prot Function]

Transcript Variant: This variant (1) utilizes an alternate 3'-terminal exon, compared to variant 2. Isoform 1 has a unique C-terminus compared to isoform 2. COMPLETENESS: complete on the 3' end.