

Product datasheet for MC208874

Ldhb (NM_008492) Mouse Untagged Clone

Product data:

Product Type:	Expression Plasmids
Product Name:	Ldhb (NM_008492) Mouse Untagged Clone
Tag:	Tag Free
Symbol:	Ldhb
Synonyms:	A1790582; H-Ld; H-Ldh; Ldh-; Ldh-2; LDH-B; LDH-H; Ldh2
Mammalian Cell Selection:	Neomycin
Vector:	pCMV6-Entry (PS100001)
E. coli Selection:	Kanamycin (25 ug/mL)
Fully Sequenced ORF:	>MC208874 representing NM_008492 Red=Cloning site Blue=ORF Orange=Stop codon

TTTGTAAATACGACTCACTATAGGGCGGCCGGAATTCGTCGACTGGATCCGGTACCGAGGAGATCTGCC
GCC**CGATCGCC**

ATGGCAACCCCTTAAGGAGAAGCTCATTGCGTCCGTTGCAGATGATGAGGCTGCCGTCCCGAACAACAAGA
TCACTGTAGTGGGCGTTGGACAAGTGGGTATGGCATGTGCCATCAGCATTCTGGGAAAGTCTCTGGCTGA
TGAACCTGGCCTGGTGGATGTGTTGGAAGACAAGCTCAAAGGAGAGATGATGGACCTGCAGCACGGGAGC
TTGTTCTCCAGACTCCGAAAATTGTGGCCGATAAAGATTACTCTGTGACAGCCAACCTCTAAGATTGTGG
TGGTGACGGCAGGAGTCCGCCAGCAGGAGGGGAGAGTCGGCTCAACCTGGTGACAGAAATGTCAACGT
GTTCAAGTTCATATTCCTCAGATCGTCAAGTACAGCCCTGACTGCACCATCATCGTGGTTTCCAACCCA
GTGGATATTCTGACTTACGTACCTGGAACCTGAGCGGGCTACCTAAGCACCGTGTGATTGGAAGCGGAT
GCAATCTGGATTCTGCTCGATTCCGCTACCTCATGGCAGAGAAGCTTGGCATTATCCCAGCAGCTGCCA
CGGATGGATCCTGGGCGAGCATGGAGACTCCAGTGTGGCTGTGTGGAGCGGGGTGAATGTGGCAGGAGTC
TCCCTCCAGGAAGTGAATCCAGAAATGGGGACAGACAATGACAGTGAGAACTGGAAGGAGGTGCATAAGA
TGGTGGTGGACAGTGCCTATGAAGTCATCAAGCTCAAAGGCTACACCAACTGGGCCATCGGCCTGAGCGT
GGCTGACCTCATCGAGTCCATGCTGAAAAACCTCTCCCGATTACCCCGTGTCTACCATGGTGAAGGGA
ATGTACGGCATTGAGAATGAAGTCTTCTCAGTCTCCCGTGCATCCTCAATGCTCGGGGGCTGACCAGCG
TCATCAATCAGAAGCTGAAGGACGATGAGGTGCTCAGCTCAGGAAGAGTGCGGACACCCGTGGGACAT
CCAGAAAGACCTCAAAGACCTGTGA

ACGCGTACGCGGCCGCTCGAGCAGAAACTCATCTCAGAAGAGGATCTGGCAGCAAATGATATCCTGGATT
ACAAGGATGACGACGATAAGGTTTAA

Chromatograms: https://cdn.origene.com/chromatograms/ja1865_c03.zip



[View online »](#)

Restriction Sites:	Sgfl-Mlul
ACCN:	NM_008492
Insert Size:	1005 bp
OTI Disclaimer:	<p>Due to the inherent nature of this plasmid, standard methods to replicate additional amounts of DNA in E. coli are highly likely to result in mutations and/or rearrangements. Therefore, OriGene does not guarantee the capability to replicate this plasmid DNA. Additional amounts of DNA can be purchased from OriGene with batch-specific, full-sequence verification at a reduced cost. Please contact our customer care team at custsupport@origene.com or by calling 301.340.3188 option 3 for pricing and delivery.</p> <p>The molecular sequence of this clone aligns with the gene accession number as a point of reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing variants is recommended prior to use. More info</p>
Components:	The ORF clone is ion-exchange column purified and shipped in a 2D barcoded Matrix tube containing 10ug of transfection-ready, dried plasmid DNA (reconstitute with 100 ul of water).
Reconstitution Method:	<ol style="list-style-type: none"> 1. Centrifuge at 5,000xg for 5min. 2. Carefully open the tube and add 100ul of sterile water to dissolve the DNA. 3. Close the tube and incubate for 10 minutes at room temperature. 4. Briefly vortex the tube and then do a quick spin (less than 5000xg) to concentrate the liquid at the bottom. 5. Store the suspended plasmid at -20°C. The DNA is stable for at least one year from date of shipping when stored at -20°C.
Note:	Plasmids are not sterile. For experiments where strict sterility is required, filtration with 0.22um filter is required.
RefSeq:	BC046755 , AAH46755
RefSeq Size:	1324 bp
RefSeq ORF:	1005 bp
Locus ID:	16832
UniProt ID:	P16125
Cytogenetics:	6 74.17 cM

Gene Summary:

This gene encodes the B subunit of lactate dehydrogenase enzyme, which catalyzes the interconversion of pyruvate and lactate with concomitant interconversion of NADH and NAD⁺ in a post-glycolysis process. Alternatively spliced transcript variants have also been found for this gene. Recent studies have shown that a C-terminally extended isoform is produced by use of an alternative in-frame translation termination codon via a stop codon readthrough mechanism, and that this isoform is localized in the peroxisomes. Pseudogenes have been identified on chromosomes 1 and 19. [provided by RefSeq, Feb 2016]

Transcript Variant: This variant (1) encodes two isoforms, which result from the use of alternative in-frame translation termination codons. The shorter isoform (Ldhb) results from translation termination at the upstream UGA stop codon, while the longer isoform (Ldhbx) results from UGA stop codon readthrough to the downstream UAG termination codon. This RefSeq represents the shorter isoform (Ldhb), which is localized in the cytosol.