

Product datasheet for MC208941

Max (NM 001146176) Mouse Untagged Clone

Product data:

Product Type: Expression Plasmids

Product Name: Max (NM_001146176) Mouse Untagged Clone

Tag: Tag Free Symbol: Max

Synonyms: AA960152; Al875693; bHLHd4; bHLHd5; bHLHd6; bHLHd7; bHLHd8

Mammalian Cell Neomycin

Selection:

Vector:pCMV6-Entry (PS100001)E. coli Selection:Kanamycin (25 ug/mL)

Fully Sequenced ORF: >MC208941 representing NM_001146176

Red=Cloning site Blue=ORF Orange=Stop codon

TTTTGTAATACGACTCACTATAGGGCGGCCGGGAATTCGTCGACTGGATCCGGTACCGAGGAGATCTGCC

GCCGCGATCGCC

CAGAGCAGGAAGAAACTCCGGATGGAGGCCAGC<mark>TAA</mark>

 ACGCGTACGCGGCCGCTCGAGCAGAAACTCATCTCAGAAGAGGATCTGGCAGCAAATGATATCCTGGATT

ACAAGGATGACGACGATAAGGTTTAA

Chromatograms: https://cdn.origene.com/chromatograms/ja3454 a04.zip

Restriction Sites: Sgfl-Mlul

ACCN: NM_001146176

Insert Size: 456 bp



OriGene Technologies, Inc. 9620 Medical Center Drive, Ste 200

CN: techsupport@origene.cn

Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com ORIGENE

OTI Disclaimer:

Our molecular clone sequence data has been matched to the reference identifier above as a point of reference. Note that the complete sequence of our molecular clones may differ from the sequence published for this corresponding reference, e.g., by representing an alternative RNA splicing form or single nucleotide polymorphism (SNP).

Components:

The ORF clone is ion-exchange column purified and shipped in a 2D barcoded Matrix tube containing 10ug of transfection-ready, dried plasmid DNA (reconstitute with 100 ul of water).

Reconstitution Method:

- 1. Centrifuge at 5,000xg for 5min.
- 2. Carefully open the tube and add 100ul of sterile water to dissolve the DNA.
- 3. Close the tube and incubate for 10 minutes at room temperature.
- 4. Briefly vortex the tube and then do a quick spin (less than 5000xg) to concentrate the liquid

at the bottom.

5. Store the suspended plasmid at -20°C. The DNA is stable for at least one year from date of

shipping when stored at -20°C.

RefSeq: <u>NM 001146176.1, NP 001139648.1</u>

 RefSeq Size:
 1978 bp

 RefSeq ORF:
 456 bp

 Locus ID:
 17187

 UniProt ID:
 P28574

Cytogenetics: 12 33.78 cM

Gene Summary: Transcription regulator. Forms a sequence-specific DNA-binding protein complex with MYC or

MAD which recognizes the core sequence 5'-CAC[GA]TG-3'. The MYC:MAX complex is a transcriptional activator, whereas the MAD:MAX complex is a repressor. CpG methylation of the recognition site greatly inhibits DNA binding, suggesting that DNA methylation may regulate the MYC:MAX complex in vivo. May repress transcription via the recruitment of a chromatin remodeling complex containing H3 'Lys-9' histone methyltransferase activity. Represses MYC transcriptional activity from E-box elements (By similarity).[UniProtKB/Swiss-

Prot Function]

based on transcript alignments.

Transcript Variant: This variant (2) lacks an exon in the 5' coding region compared to variant 1 but maintains the reading frame. The encoded protein (isoform 2) is shorter but has the same N- and C-termini compared to isoform 1. Sequence Note: This RefSeq record was created from transcript and genomic sequence data to make the sequence consistent with the reference genome assembly. The genomic coordinates used for the transcript record were