

Product datasheet for **MC209178**

Pkia (NM_008862) Mouse Untagged Clone

Product data:

Product Type:	Expression Plasmids
Product Name:	Pkia (NM_008862) Mouse Untagged Clone
Tag:	Tag Free
Symbol:	Pkia
Synonyms:	AI415001; PKIalpha
Mammalian Cell Selection:	Neomycin
Vector:	pCMV6-Entry (PS100001)
E. coli Selection:	Kanamycin (25 ug/mL)
Fully Sequenced ORF:	>NCBI ORF sequence for NM_008862, the custom clone sequence may differ by one or more nucleotides

```
ATGACTGATGTGGAACTACGTATGCAGATTTTCATTGCTTCAGGAAGAACAGGTAGAAGAAATGCAATAC
ATGATATCCTGGTTTCCTCTGCAAGTGGCAACAGCAATGAATTAGCCTTAAAACTAGCAGGCCTTGATAT
CAACAAGACAGAAGGTGAAGATGATGGACAGAGAAGCTCCACCGAACAAAGTGGAGAAGCCAGGGAGAA
GCAGCCAAGTCTGAAAGCTAA
```

Restriction Sites:	Sgfl-MluI
ACCN:	NM_008862
Insert Size:	231 bp
OTI Disclaimer:	Our molecular clone sequence data has been matched to the reference identifier above as a point of reference. Note that the complete sequence of our molecular clones may differ from the sequence published for this corresponding reference, e.g., by representing an alternative RNA splicing form or single nucleotide polymorphism (SNP).
Components:	The ORF clone is ion-exchange column purified and shipped in a 2D barcoded Matrix tube containing 10ug of transfection-ready, dried plasmid DNA (reconstitute with 100 ul of water).



[View online »](#)

Reconstitution Method:

1. Centrifuge at 5,000xg for 5min.
2. Carefully open the tube and add 100ul of sterile water to dissolve the DNA.
3. Close the tube and incubate for 10 minutes at room temperature.
4. Briefly vortex the tube and then do a quick spin (less than 5000xg) to concentrate the liquid at the bottom.
5. Store the suspended plasmid at -20°C. The DNA is stable for at least one year from date of shipping when stored at -20°C.

RefSeq: [BC048244](#), [AAH48244](#)

RefSeq Size: 3640 bp

RefSeq ORF: 231 bp

Locus ID: 18767

UniProt ID: [P63248](#)

Cytogenetics: 3 A1

Gene Summary: Extremely potent competitive inhibitor of cAMP-dependent protein kinase activity, this protein interacts with the catalytic subunit of the enzyme after the cAMP-induced dissociation of its regulatory chains.[UniProtKB/Swiss-Prot Function]