

Product datasheet for **MC209225**

Prkag1 (NM_016781) Mouse Untagged Clone

Product data:

Product Type: Expression Plasmids
Product Name: Prkag1 (NM_016781) Mouse Untagged Clone
Tag: Tag Free
Symbol: Prkag1
Synonyms: AA571379; BB036179; Prkaac
Vector: pCMV6-Entry (PS100001)
E. coli Selection: Kanamycin (25 ug/mL)
Cell Selection: Neomycin
Fully Sequenced ORF: >MC209225 representing NM_016781
Red=Cloning site Blue=ORF Orange=Stop codon

TTTTGTAATACGACTCACTATAGGGCGCCGGAATTCGTCGACTGGATCCGGTACCGAGGAGATCTGCC
GCC**GCGATCGCC**

ATGGAGTCGGTTGCTGCAGAGAGCTCGCCAGCTCTAGAGAATGAACACTTTC AAGAGACCCCGAATCAA
ACAATAGTGTATACTTCCTTCATGAAGTCTCATCGCTGCTATGACCTAATCCACAAGTTCCAAGTT
GGTGGTATTTGACTTTCGCTACAGGTAAGAAAGCCCTTTTTGTCCTGGTGACCAATGGTGTTCGTGCC
GCCCTTTGTGGGACAGTAAGAAGCAGAGTTTTGTGGCATGCTGACCATCACCGACTTCATCAACATTT
TGCACCGATACTATAAGTCAGCCCTGGTGCAGATCTACGAACTGGAGGAGCACAAGATAGAGACGTGGAG
AGAGGTGTACCTGCAGGACTCCTTTAAGCCACTTGTCTGCATCTCTCAAATGCCAGCTTGTGGTGTCT
GTCTCTTCATTAATTCGAAATAAGATCCACAGGCTCCCAGTTATCGACCCAGAGTCAGGCAACACCTTGT
ACATCCTTACTCACAAGCGGATCCTCAAGTTCCTCAAGTTGTTTATCACCGAGTCCCAAGCCGGAATT
CATGTCTAAGTCTCTCAAGAGCTGCAGATTGGCACCTATGCCAATATTGCCATGGTCCGTAACACAG
CCTGTCTACGTGGCTCTGGGCATCTTTGTACAGCACCGAGTCTCCGCCTTACCTGTAGTGGATGAGAAAG
GGCGTGTGGTGGACATCTACTCAAGTTTGTATGTGATCAATTTGGCAGCCGAAAGACCTACAACAACCT
AGATGTGTCTGTGACAAAAGCCCTGCAGCATCGGTCCCCTACTTTGAGGGTGTCTCAAATGCTACCTG
CATGAGACTCTGAAAACCATCATCAATAGGCTGGTGGAGGCAGAGGTTACCCGTCTGGTGGTGGTGGATG
AACACGACGTGGTCAAGGGCATCGTTTCGCTGTCTGACATCTTACAGGCTCTGGTGTCTACGGTGGAGA
GAAGAAGCC**TGA**

ACGCGTACGCGGCCGCTCGAGCAGAACTCATCTCAGAAGAGGATCTGGCAGCAATGATATCCTGGATT
ACAAGGATGACGACGATAAGGTTTAA

Restriction Sites: SgfI-MluI
ACCN: NM_016781



[View online »](#)

Insert Size: 993 bp

OTI Disclaimer: Due to the inherent nature of this plasmid, standard methods to replicate additional amounts of DNA in E. coli are highly likely to result in mutations and/or rearrangements. Therefore, OriGene does not guarantee the capability to replicate this plasmid DNA. Additional amounts of DNA can be purchased from OriGene with batch-specific, full-sequence verification at a reduced cost. Please contact our customer care team at custsupport@origene.com or by calling 301.340.3188 option 3 for pricing and delivery.

The molecular sequence of this clone aligns with the gene accession number as a point of reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing variants is recommended prior to use. [More info](#)

Components: The ORF clone is ion-exchange column purified and shipped in a 2D barcoded Matrix tube containing 10ug of transfection-ready, dried plasmid DNA (reconstitute with 100 ul of water).

Reconstitution Method:

1. Centrifuge at 5,000xg for 5min.
2. Carefully open the tube and add 100ul of sterile water to dissolve the DNA.
3. Close the tube and incubate for 10 minutes at room temperature.
4. Briefly vortex the tube and then do a quick spin (less than 5000xg) to concentrate the liquid at the bottom.
5. Store the suspended plasmid at -20°C. The DNA is stable for at least one year from date of shipping when stored at -20°C.

RefSeq: [NM_016781.2](#), [NP_058061.2](#)

RefSeq Size: 1680 bp

RefSeq ORF: 993 bp

Locus ID: 19082

UniProt ID: [O54950](#)

Cytogenetics: 15 54.73 cM

Gene Summary:

AMP/ATP-binding subunit of AMP-activated protein kinase (AMPK), an energy sensor protein kinase that plays a key role in regulating cellular energy metabolism. In response to reduction of intracellular ATP levels, AMPK activates energy-producing pathways and inhibits energy-consuming processes: inhibits protein, carbohydrate and lipid biosynthesis, as well as cell growth and proliferation. AMPK acts via direct phosphorylation of metabolic enzymes, and by longer-term effects via phosphorylation of transcription regulators. Also acts as a regulator of cellular polarity by remodeling the actin cytoskeleton; probably by indirectly activating myosin. Gamma non-catalytic subunit mediates binding to AMP, ADP and ATP, leading to activate or inhibit AMPK: AMP-binding results in allosteric activation of alpha catalytic subunit (PRKAA1 or PRKAA2) both by inducing phosphorylation and preventing dephosphorylation of catalytic subunits. ADP also stimulates phosphorylation, without stimulating already phosphorylated catalytic subunit. ATP promotes dephosphorylation of catalytic subunit, rendering the AMPK enzyme inactive (By similarity).[UniProtKB/Swiss-Prot Function]