

## Product datasheet for MC209262

## Sirpa (NM\_001177647) Mouse Untagged Clone

## **Product data:**

## OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

Product Type:	Expression Plasmids
Product Name:	Sirpa (NM_001177647) Mouse Untagged Clone
Tag:	Tag Free
Symbol:	Sirpa
Synonyms:	Al835480; Bit; CD172a; P84; Ptpns1; SHP-1; SHPS-1; SIRP
Vector:	pCMV6-Entry (PS100001)
E. coli Selection:	Kanamycin (25 ug/mL)
Cell Selection:	Neomycin
Fully Sequenced ORF:	>MC209262 representing NM_001177647 Red=Cloning site Blue=ORF Orange=Stop codon
	TTTTGTAATACGACTCACTATAGGGCGGCCGGGAATTCGTCGACTGGATCCGGTACCGAGGAGATCTGCC GCC <mark>GCGATCGC</mark> C
	ATGGAGCCCGGCCGGCCCGGGCCCTGGCCGCTAGGGCCGCTGCTGCTGCTGCTGCTGCTCCCGCGTCCT GTTTCTGTACAGGAGCCACGGGGAAGGAACTGAAGGTGACTCAGCCTGAGAAATCAGTGTCTGTTGCTGC TGGGGATTCGACCGTTCTGAACTGCACTTTGACCTCCTTGTTGCCGGTGGGACCCATTAGGTGGTACAGA GGAGTAGGGCCAAGCCGGCTGTTGATCTACAGTTTCGCAGGAGAATACGTTCCTCGAATTAGAAATGTTT CAGATACTACTAAGAGAAACAATATGGACTTTTCCATCCGTATCAGTAATGTCACCCCAGCAGATGCTGG CATCTACTACTGTGTGAAGTTCCAGAAAGGATCATCAGAGCCTGACACAGAAATACAATCTGGAGGGGGA ACAGAGGTCTATGTACTCGATAATAATGCTACCACAACTGGAATGTCTTCATCGGTGTGGGCGTGGCGT GTGCTTTGCTCGTAGTCCTGCTGATGGCTGCTCTCTACCTCCTCGGATCAAACAGAAGAAAGCCAAGGG GTCAACATCTTCCACACGGTTGCACGAGCCCGAGAAGAACGCCAGGGAAATAACCCAGAACACA AATGACATCAACGACATCACATACGCAGACCTGAATCTGCCCAAAGAGAAGACGCCAGACACA AATGACATCAACGACATCACATACGCAGACCTGAATCTGCCCAAAGGAGAAGAAGCCCGCACCCGGGCCC CTGAGCCTAACAACCACACAGAATATGCAAGCATTGAGACAGGCAAGGCCAGGCCAAGGCCAGGGAAATACCCT CACCTATGCTGGACATGGTCCACCTCAGCCGGGCACAGCCAGGCCAAGCCCGAGCCAGGGATACCCT CACCTATGCTGACCTGGACATGGTCCACCTCAGCCGGGCACAGCCAGGCCAAGCCCGAGCCAGGCAATTGCCAGGCCAGGCCCGAGGAAGTGG ACGCGTACGCGCCGCTCGAGCAGGAAACTCCATCTCCAGGACGAGGCAAAGGCCCGCACCTGAGCCAGGCCACGCCCCAAGCCTGAGCCAGGCCCCAAGCCTGAGCCAGGCCACAGCCAGGCAAATGGCCAGGCCAACTCTTCC TCAGAGTATGCTAGTGTCCAGGTCCAGAGAAACTCCATCTCCAGGCCAGGCAAATGGCCAGGCAAATGGTCCAGGCAGG
Destriction Citere	
Restriction Sites:	Sgfl-Mlul
ACCN:	NM_001177647
Insert Size:	876 bp



This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2022 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US

Sirpa (NM_001177647) Mouse Untagged Clone – MC209262	
OTI Disclaimer:	Our molecular clone sequence data has been matched to the reference identifier above as a point of reference. Note that the complete sequence of our molecular clones may differ from the sequence published for this corresponding reference, e.g., by representing an alternative RNA splicing form or single nucleotide polymorphism (SNP).
Components:	The ORF clone is ion-exchange column purified and shipped in a 2D barcoded Matrix tube containing 10ug of transfection-ready, dried plasmid DNA (reconstitute with 100 ul of water).
Reconstitution Method:	<ol> <li>Centrifuge at 5,000xg for 5min.</li> <li>Carefully open the tube and add 100ul of sterile water to dissolve the DNA.</li> <li>Close the tube and incubate for 10 minutes at room temperature.</li> <li>Briefly vortex the tube and then do a quick spin (less than 5000xg) to concentrate the liquid at the bottom.</li> <li>Store the suspended plasmid at -20°C. The DNA is stable for at least one year from date of shipping when stored at -20°C.</li> </ol>
RefSeq:	<u>NM 001177647.2, NP 001171118.1</u>
RefSeq Size:	3377 bp
RefSeq ORF:	876 bp
Locus ID:	19261
UniProt ID:	<u>P97797</u>
Cytogenetics:	2 63.19 cM
Gene Summary:	Immunoglobulin-like cell surface receptor for CD47. Acts as docking protein and induces translocation of PTPN6, PTPN11 and other binding partners from the cytosol to the plasma membrane. Supports adhesion of cerebellar neurons, neurite outgrowth and glial cell attachment. May play a key role in intracellular signaling during synaptogenesis and in synaptic function. Involved in the negative regulation of receptor tyrosine kinase-coupled cellular responses induced by cell adhesion, growth factors or insulin. Mediates negative regulation of phagocytosis, mast cell activation and dendritic cell activation. CD47 binding prevents maturation of immature dendritic cells and inhibits cytokine production by mature dendritic cells (By similarity).[UniProtKB/Swiss-Prot Function] Transcript Variant: This variant (3) lacks two alternate in-frame exons in the central coding region, and uses an alternate in-frame splice site in the 3' coding region, compared to variant 4. The encoded isoform (2) is shorter than isoform 3.

This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2022 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US