

## **Product datasheet for MC209298**

## 9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

OriGene Technologies, Inc.

## Rbbp6 (NM\_175023) Mouse Untagged Clone

**Product data:** 

**Product Type:** Expression Plasmids

Product Name: Rbbp6 (NM\_175023) Mouse Untagged Clone

Tag: Tag Free Symbol: Rbbp6

**Synonyms:** 4933422O15Rik; Al316869; BB233631; C030034J04Rik; C77662; P2P-R; PACT

Mammalian Cell

Selection:

Insert Size:

Neomycin

**Vector:** pCMV6-Entry (PS100001) **E. coli Selection:** Kanamycin (25 ug/mL)

Fully Sequenced ORF: >MC209298 representing NM\_175023

372 bp

Red=Cloning site Blue=ORF Orange=Stop codon

TTTTGTAATACGACTCACTATAGGGCGGCCGGGAATTCGTCGACTGGATCCGGTACCGAGGAGATCTGCC

GCCGCGATCGCC

TTATAATGTTTCAGTGACGTAA

**ACGCGT**ACGCGGCCGCTCGAGCAGAAACTCATCTCAGAAGAGGATCTGGCAGCAAATGATATCCTGGATT

ACAAGGATGACGACGATAAGGTTTAA

**Restriction Sites:** Sgfl-Mlul ACCN: NM\_175023

**OTI Disclaimer:** Our molecular clone sequence data has been matched to the reference identifier above as a

point of reference. Note that the complete sequence of our molecular clones may differ from the sequence published for this corresponding reference, e.g., by representing an alternative

RNA splicing form or single nucleotide polymorphism (SNP).





Components:

The ORF clone is ion-exchange column purified and shipped in a 2D barcoded Matrix tube containing 10ug of transfection-ready, dried plasmid DNA (reconstitute with 100 ul of water).

**Reconstitution Method:** 

- 1. Centrifuge at 5,000xg for 5min.
- 2. Carefully open the tube and add 100ul of sterile water to dissolve the DNA.
- 3. Close the tube and incubate for 10 minutes at room temperature.
- 4. Briefly vortex the tube and then do a quick spin (less than 5000xg) to concentrate the liquid

at the bottom.

5. Store the suspended plasmid at -20°C. The DNA is stable for at least one year from date of

shipping when stored at -20°C.

**Note:** Plasmids are not sterile. For experiments where strict sterility is required, filtration with

0.22um filter is required.

RefSeq: <u>NM 175023.3</u>, <u>NP 778188.1</u>

 RefSeq Size:
 2272 bp

 RefSeq ORF:
 372 bp

 Locus ID:
 19647

 UniProt ID:
 P97868

 Cytogenetics:
 7 F2

**Gene Summary:** E3 ubiquitin-protein ligase which promotes ubiquitination of YBX1, leading to its degradation

by the proteasome (By similarity). May play a role as a scaffold protein to promote the assembly of the p53/TP53-MDM2 complex, resulting in increase of MDM2-mediated

ubiquitination and degradation of p53/TP53; may function as negative regulator of p53/TP53, leading to both apoptosis and cell growth retardation (PubMed:17470788). Regulates DNA-replication and common fragile sites (CFS) stability in a ZBTB38- and MCM10-dependent

manner. Controls ZBTB38 protein stability and abundance via ubiquitination and

proteasomal degradation, and ZBTB38 in turn negatively regulates the expression of MCM10 which plays an important role in DNA-replication (PubMed:24726359).[UniProtKB/Swiss-Prot

Function]

Transcript Variant: This variant (2) lacks most 3' exons and has an alternate 3' segment, as

compared to variant 1. The resulting isoform 2 is much shorter than isoform 1.