

Product datasheet for **MC209378**

Sell (NM_001164059) Mouse Untagged Clone

Product data:

Product Type:	Expression Plasmids
Product Name:	Sell (NM_001164059) Mouse Untagged Clone
Tag:	Tag Free
Symbol:	Sell
Synonyms:	AI528707; CD62L; L-selectin; LECAM-1; Lnhrr; Ly-22; Ly-m22; Lyam-1; Lyam1
Mammalian Cell Selection:	Neomycin
Vector:	pCMV6-Entry (PS100001)
E. coli Selection:	Kanamycin (25 ug/mL)
Restriction Sites:	Sgfl-Mlul
ACCN:	NM_001164059
Insert Size:	1011 bp
OTI Disclaimer:	Our molecular clone sequence data has been matched to the reference identifier above as a point of reference. Note that the complete sequence of our molecular clones may differ from the sequence published for this corresponding reference, e.g., by representing an alternative RNA splicing form or single nucleotide polymorphism (SNP).
Components:	The ORF clone is ion-exchange column purified and shipped in a 2D barcoded Matrix tube containing 10ug of transfection-ready, dried plasmid DNA (reconstitute with 100 ul of water).
Reconstitution Method:	<ol style="list-style-type: none"> 1. Centrifuge at 5,000xg for 5min. 2. Carefully open the tube and add 100ul of sterile water to dissolve the DNA. 3. Close the tube and incubate for 10 minutes at room temperature. 4. Briefly vortex the tube and then do a quick spin (less than 5000xg) to concentrate the liquid at the bottom. 5. Store the suspended plasmid at -20°C. The DNA is stable for at least one year from date of shipping when stored at -20°C.
RefSeq:	<u>NM_001164059.1, NP_001157531.1</u>
RefSeq Size:	2213 bp
RefSeq ORF:	1011 bp


[View online »](#)

Locus ID: 20343

Cytogenetics: 1 71.37 cM

Gene Summary: Calcium-dependent lectin that mediates cell adhesion by binding to glycoproteins on neighboring cells. Mediates the adherence of lymphocytes to endothelial cells of high endothelial venules in peripheral lymph nodes (PubMed:1693096). Promotes initial tethering and rolling of leukocytes in endothelia (By similarity).[UniProtKB/Swiss-Prot Function]
Transcript Variant: This variant (2) lacks an alternate in-frame exon in the central coding region, compared to variant 1. The resulting isoform (2) lacks an internal segment, compared to isoform 1. Sequence Note: The RefSeq transcript and protein were derived from genomic sequence to make the sequence consistent with the reference genome assembly. The genomic coordinates used for the transcript record were based on alignments.