

## Product datasheet for **MC209507**

### **Tdg (NM\_172552) Mouse Untagged Clone**

#### **Product data:**

Product Type:	Expression Plasmids
Product Name:	Tdg (NM_172552) Mouse Untagged Clone
Tag:	Tag Free
Symbol:	Tdg
Synonyms:	E130317C12Rik; JZA-3; Jza1
Mammalian Cell Selection:	Neomycin
Vector:	pCMV6-Entry (PS100001)
E. coli Selection:	Kanamycin (25 ug/mL)



[View online »](#)

**Fully Sequenced ORF:** >MC209507 representing NM\_172552  
 Red=Cloning site Blue=ORF Orange=Stop codon

TTTTGTAATACGACTCACTATAGGGCGGCCGGAATTCGTCGACTGGATCCGGTACCGAGGAGATCTGCC  
 GCC**CGGATCGC**

ATGGACGCAGAGGCCGCGCAGCTATTCTCTGGAGCAAGTTCAAGCTTTGTATTTCATTTCATTCAAC  
 AAATGATGGCAGAAGTTCCTAACATGGCAGTCACGACTGGACAGCAGGTGCCAGCAGTAGCTCCTAACAT  
 GGCAACCGTGACAGAACAGCAGGTGCCGAAGACGCTCCTGTCCAGGAACCTGCACCAGAAGCTCCAAAG  
 AGAAGGAAAAGGAAACCCAGAGCAGCAGAGCCCCAGGAACAGTGGAGCCCAAAAAACCTGCTACGTCGA  
 AGAAATCCGGCAAGTCTACAAAATCAAAGGAAAAGCAGGAGAAAATCACAGACGCGTTTAAAGTAAAAG  
 GAAAGTGGACCGCTTCAACGGCGTCTCTGAAGCTGAGCTTCTGACCAAGACTCTTCCTGACATTTTGACC  
 TTCAATCTGGATATTGTGATCATTGGCATTAAACCGGGATTAATGGCTGCTTACAAGGACATCACTACC  
 CTGGGCTGGAAATCACTTCTGGAAGTGTCTGTTCATGTCGGGGCTGAGTGAGGTGCAGCTGAATCACAT  
 GGATGACCACACCTTACCGGCAAGTACGGCATCGGATTCACCAACATGGTGGAACGGACGACCCGGGC  
 AGCAAGGATCTGTCTAGTAAAGAGTTCCGGGAAGGAGGGCGCATCCTGGTGCAGAACTGCAGAAATATC  
 AGCCACGAATAGCGGTGTTTAAATGGAATGTATTTATGAAATTTTCAAGTAAAGAGTTTTTGGAGTAA  
 GGTTAAGAACTTGAATTTGGGCTTCAACCCACAAGATCCCAGACACAGAACTCTGTGCTACGTCATG  
 CCGTCGTCCAGCGCCAGATGTGCTCAGTTTCCCGGGCCAGGACAAAGTTCATTACTACATTAAGCTGA  
 AGGACTTGAGAGACCAACTGAAAGGCATTGAACGCAACGCGGACGTTTCAAGGAAGTGCAGTATACATTGA  
 CCTGCAGCTTGCGCAAGAGGACGCAAGAAGATGGCTGTTAAGGAAGAAAAGTATGATCCAGGCTATGAG  
 GCAGCTTACGGCGGTGCCTATGGGAAAACCATGTAAATGGGGAACCTTGTGGCATTGCTTCAAATGGGC  
 TAACAGCTCACAGTGGGAGCCGAGAGGAGAAGCGGCCCCAGCGATGTTCCGAATGGGAGTGGATGGC  
 ACAGTCGTTTGCAGAGCAGATCCCTTCTTTAATAATTGTGGGACCCGAGAGCAGGAAGAAGAGAGCCAC  
 GCT**AG**

AG**CGGACCG**ACGCGTACGCGCCGCTCGAGCAGAACTCATCTCAGAAGAGGATCTGGCAGCAAATGATATCC  
 TGGATTACAAGGATGACGACGATAAGGTTTAA

**Restriction Sites:** SgfI-RsrII

**ACCN:** NM\_172552

**Insert Size:** 1266 bp

**OTI Disclaimer:** Our molecular clone sequence data has been matched to the reference identifier above as a point of reference. Note that the complete sequence of our molecular clones may differ from the sequence published for this corresponding reference, e.g., by representing an alternative RNA splicing form or single nucleotide polymorphism (SNP).

**Components:** The ORF clone is ion-exchange column purified and shipped in a 2D barcoded Matrix tube containing 10ug of transfection-ready, dried plasmid DNA (reconstitute with 100 ul of water).

**Reconstitution Method:**

1. Centrifuge at 5,000xg for 5min.
2. Carefully open the tube and add 100ul of sterile water to dissolve the DNA.
3. Close the tube and incubate for 10 minutes at room temperature.
4. Briefly vortex the tube and then do a quick spin (less than 5000xg) to concentrate the liquid at the bottom.
5. Store the suspended plasmid at -20°C. The DNA is stable for at least one year from date of shipping when stored at -20°C.

<b>Note:</b>	Plasmids are not sterile. For experiments where strict sterility is required, filtration with 0.22um filter is required.
<b>RefSeq:</b>	<a href="#">NM_172552.3</a> , <a href="#">NP_766140.2</a>
<b>RefSeq Size:</b>	3218 bp
<b>RefSeq ORF:</b>	1266 bp
<b>Locus ID:</b>	21665
<b>UniProt ID:</b>	<a href="#">P56581</a>
<b>Cytogenetics:</b>	10 39.72 cM
<b>Gene Summary:</b>	<p>DNA glycosylase that plays a key role in active DNA demethylation: specifically recognizes and binds 5-formylcytosine (5fC) and 5-carboxylcytosine (5caC) in the context of CpG sites and mediates their excision through base-excision repair (BER) to install an unmethylated cytosine (PubMed:21817016). Cannot remove 5-hydroxymethylcytosine (5hmC). According to an alternative model, involved in DNA demethylation by mediating DNA glycolase activity toward 5-hydroxymethyluracil (5hmU) produced by deamination of 5hmC (PubMed:21722948). Also involved in DNA repair by acting as a thymine-DNA glycosylase that mediates correction of G/T mispairs to G/C pairs: in the DNA of higher eukaryotes, hydrolytic deamination of 5-methylcytosine to thymine leads to the formation of G/T mismatches. Its role in the repair of canonical base damage is however minor compared to its role in DNA demethylation. It is capable of hydrolyzing the carbon-nitrogen bond between the sugar-phosphate backbone of the DNA and a mispaired thymine. In addition to the G/T, it can remove thymine also from C/T and T/T mispairs in the order G/T &gt;&gt; C/T &gt; T/T. It has no detectable activity on apyrimidinic sites and does not catalyze the removal of thymine from A/T pairs or from single-stranded DNA. It can also remove uracil and 5-bromouracil from mispairs with guanine. [UniProtKB/Swiss-Prot Function]</p> <p>Transcript Variant: This variant (3) represents the longer variant and encodes the longest isoform (2).</p>