

Product datasheet for MC209750

Macroh2a1 (NM 001159513) Mouse Untagged Clone

Product data:

Product Type: Expression Plasmids

Product Name: Macroh2a1 (NM_001159513) Mouse Untagged Clone

Tag: Tag Free

Symbol: Macroh2a1

Synonyms: H2af; H2AF12; H2AF12M; H2afy; MACROH2; mH2a; mH2a1

Vector:pCMV6-Entry (PS100001)E. coli Selection:Kanamycin (25 ug/mL)

Cell Selection: Neomycin

Fully Sequenced ORF: >MC209750 representing NM_001159513

Red=Cloning site Blue=ORF Orange=Stop codon

TTTTGTAATACGACTCACTATAGGGCGGCCGGGAATTCGTCGACTGGATCCGGTACCGAGGAGATCTGCC

GCCGCGATCGCC

ATGTCGAGCCGCGGGGAAGAAGAAATCCACCAAGACCTCCCGGTCAGCCAAGGCCGGAGTCATCTTCC CTGTGGGACGCATGCTTCGGTACATCAAGAAAGGCCACCCTAAGTATAGGATCGGAGTGGGGGCACCTGT AACAAGAAGGGACGGTCACACCCCGGCACATCCTGTTAGCTGTGGCCAATGATGAAGAGCTAAACCAGC GCGAGGATCCAAGGGAAAATTGGAAGCCATCATCACGCCTCCGCCGGCCAAAAAAGGCCAAGTCTCCATCC CAGAAGAAGCCAGTGGCTAAGAAGACAGGAGGCAAGAAAGGGGCCCGGAAGTCTAAGAAGCAGGGAGAAG TGAGCAAGGCGGCCAGCGCAGACAGTACGACGGAGGGCACGCCTACAGACGGCTTCACTGTCCTCTCCAC CAAGAGCCTCTTCCTCGGCCAGAAGCTGAACCTTATTCACAGTGAAATCAGTAATTTAGCCGGCTTTGAG GTGGAGGCCATAATCAATCCTACCAATGCTGACATTGACCTTAAAGATGACCTAGGAAACACACTGGAGA AGAAGGGCGGCAAGGAGTTTGTAGAAGCTGTTCTGGAACTCCGGAAAAAGAACGGGCCCTTGGAGGTAGC TGGAGCTGCTATTAGTGCAGGCCATGGCCTGCCAAGTTTGTGATCCACTGTAATAGTCCTGTCTGG GGTGCAGACAAATGTGAAGAACTTCTAGAAAAGACGGTGAAAAACTGCTTGGCTCTAGCTGATGACAGAA AGCTGAAATCCATCGCCTTCCCATCCATTGGCAGCGGCAGGAACGGGTTCCCGAAGCAGACAGCGGCCCA GCTCATTCTGAAGGCCATCTCCAGCTACTTTGTCTCCACGATGTCCTCCTCCATCAAAACTGTGTACTTC ATGCTTTTTGACAGTGAGAGCATAGGTATCTATGTGCAGGAAATGGCCAAGCTGGACGCCAAC<mark>TAG</mark>

ACGCGTACGCGGCCGCTCGAGCAGAAACTCATCTCAGAAGAGGATCTGGCAGCAAATGATATCCTGGATT

ACAAGGATGACGACGATAAGGTTTAA

Restriction Sites: Sgfl-Mlul

ACCN: NM_001159513



OriGene Technologies, Inc. 9620 Medical Center Drive, Ste 200

CN: techsupport@origene.cn

Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com ORIGENE

Insert Size: 1116 bp

OTI Disclaimer: Our molecular clone sequence data has been matched to the reference identifier above as a

point of reference. Note that the complete sequence of our molecular clones may differ from the sequence published for this corresponding reference, e.g., by representing an alternative

RNA splicing form or single nucleotide polymorphism (SNP).

Components: The ORF clone is ion-exchange column purified and shipped in a 2D barcoded Matrix tube

containing 10ug of transfection-ready, dried plasmid DNA (reconstitute with 100 ul of water).

Reconstitution Method: 1. Centrifuge at 5,000xg for 5min.

2. Carefully open the tube and add 100ul of sterile water to dissolve the DNA.

3. Close the tube and incubate for 10 minutes at room temperature.

4. Briefly vortex the tube and then do a quick spin (less than 5000xg) to concentrate the liquid

at the bottom.

5. Store the suspended plasmid at -20°C. The DNA is stable for at least one year from date of

shipping when stored at -20°C.

RefSeq: <u>NM 001159513.1</u>, <u>NP 001152985.1</u>

13 B1

 RefSeq Size:
 1975 bp

 RefSeq ORF:
 1116 bp

 Locus ID:
 26914

 UniProt ID:
 Q9QZQ8

Cytogenetics:

Gene Summary: Histones are basic nuclear proteins that are responsible for the nucleosome structure of the

chromosomal fiber in eukaryotes. Nucleosomes consist of approximately 146 bp of DNA wrapped around a histone octamer composed of pairs of each of the four core histones (H2A, H2B, H3, and H4). The chromatin fiber is further compacted through the interaction of a linker histone, H1, with the DNA between the nucleosomes to form higher order chromatin structures. This gene encodes a replication-independent histone that is a member of the histone H2A family. It replaces conventional H2A histones in a subset of nucleosomes where it represses transcription and participates in stable X chromosome inactivation. Alternative splicing results in multiple transcript variants encoding different isoforms. [provided by

RefSeq, Nov 2015]

Transcript Variant: This variant (2) uses an alternate in-frame splice site in the central coding region, compared to variant 1. The resulting isoform (2) lacks one internal residue, compared

to isoform 1.