

Product datasheet for **MC209860**

Sirt6 (NM_001163430) Mouse Untagged Clone

Product data:

Product Type: Expression Plasmids
Product Name: Sirt6 (NM_001163430) Mouse Untagged Clone
Tag: Tag Free
Symbol: Sirt6
Synonyms: 2810449N18Rik; AI043036; Sir2l6
Vector: pCMV6-Entry (PS100001)
E. coli Selection: Kanamycin (25 ug/mL)
Cell Selection: Neomycin
Fully Sequenced ORF: >MC209860 representing NM_001163430
 Red=Cloning site Blue=ORF Orange=Stop codon

TTTTGTAATACGACTCACTATAGGGCGCCGGGAATTCGTCGACTGGATCCGGTACCGAGGAGATCTGCC
 GCC**GCGATCGCC**

ATGTGGCAGTCTCCAGCGTGGTTTTCCACACGGGCGCCGGCATCAGCACCGCCTCTGGCATCCCCGACT
 TCAGAGGCCCCCATGGCGTGTGGACCATGGAGGAACGCGGCTGGCCCCAAGTTTGACACCACCTTCGA
 GAATGCTCGGCCCTCGAAGACCCACATGGCCCTGGTTCAGCTAGAACGCATGGGCTTCTCAGCTTCTG
 GTAGCCAGAACGTAGACGGGCTGCACGTGCGCTCGGGCTTCCCCAGGGACAAGCTGGCAGAGCTGCACG
 GAAACATGTTGTAGAGGAATGTCCCAAGTGTAAAGACGCAGTACGTGAGACACGGTTGTGGGCACCAT
 GGGCCTCAAGGCCACAGGCCGGCTCTGCACCGTGGCCAAGACCAGGGGACTTCGGGCCTGTAGAGGGGAG
 CTGAGAGACACCATTTCTGGACTGGGAGGACTCGTTGCCTGACCGGGACCTGATGCTCGCTGATGAGGCCA
 GCAGGACCGCAGACCTGTCTGTCAACCCTGGGTACCTCGCTGCAGATCCGCCCCAGTGGGAACCTGCCCT
 TGCCACTAAGCGCCGAGGAGGCCGTCTGGTCATTGTCAACCTGCAACCCACAAAACATGACCGCCAGGT
 GACCTGCGCATCCACGGCTACGTGGATGAGGTGATGTGCAGACTCATGAAGCATCTGGGGCTGGAGATTC
 CAGCCTGGGATGGACCCGCGTGTAGACAAAGCCCTGCCACCTCGCTCGCCAGTAGCACTCAAGGC
 TGAGCCCCCGTGCATCTCAATGGTGCAGTGCATGTTTCGTATAAGTCCAAGCCCAACAGCCCTATACTC
 CACAGGCCCCCAAAAGAGTGAAGACCGAGGCTGCCCCAGCTGA

ACGCGTACGCGGCCGCTCGAGCAGAAACTCATCTCAGAAGAGGATCTGGCAGCAAATGATATCCTGGATT
 ACAAGGATGACGACGATAAGGTTTAA

Restriction Sites: SgfI-MluI
ACCN: NM_001163430
Insert Size: 885 bp



[View online »](#)

OTI Disclaimer:	Our molecular clone sequence data has been matched to the reference identifier above as a point of reference. Note that the complete sequence of our molecular clones may differ from the sequence published for this corresponding reference, e.g., by representing an alternative RNA splicing form or single nucleotide polymorphism (SNP).
Components:	The ORF clone is ion-exchange column purified and shipped in a 2D barcoded Matrix tube containing 10ug of transfection-ready, dried plasmid DNA (reconstitute with 100 ul of water).
Reconstitution Method:	<ol style="list-style-type: none">1. Centrifuge at 5,000xg for 5min.2. Carefully open the tube and add 100ul of sterile water to dissolve the DNA.3. Close the tube and incubate for 10 minutes at room temperature.4. Briefly vortex the tube and then do a quick spin (less than 5000xg) to concentrate the liquid at the bottom.5. Store the suspended plasmid at -20°C. The DNA is stable for at least one year from date of shipping when stored at -20°C.
RefSeq:	<u>NM_001163430.1, NP_001156902.1</u>
RefSeq Size:	1668 bp
RefSeq ORF:	885 bp
Locus ID:	50721
Cytogenetics:	10 39.72 cM
Gene Summary:	<p>NAD-dependent protein deacetylase. Has deacetylase activity towards histone H3K9Ac and H3K56Ac. Modulates acetylation of histone H3 in telomeric chromatin during the S-phase of the cell cycle. Deacetylates histone H3K9Ac at NF-kappa-B target promoters and may down-regulate the expression of a subset of NF-kappa-B target genes. Deacetylation of nucleosomes interferes with RELA binding to target DNA. May be required for the association of WRN with telomeres during S-phase and for normal telomere maintenance. On DNA damage, promotes DNA end resection via deacetylation of RBBP8. Has very weak deacetylase activity and can bind NAD(+) in the absence of acetylated substrate (By similarity). Acts as a corepressor of the transcription factor Hif1a to control the expression of multiple glycolytic genes to regulate glucose homeostasis. Required for genomic stability. Required for normal IGF1 serum levels and normal glucose homeostasis. Modulates cellular senescence and apoptosis. Regulates the production of TNF protein. Has a role in the regulation of life span in male mice, but not in female mice.[UniProtKB/Swiss-Prot Function]</p> <p>Transcript Variant: This variant (2) differs in the UTRs and 5' coding region and initiates translation at a downstream start codon, compared to variant 1. The resulting isoform (2) has a shorter N-terminus compared to isoform 1.</p>