

## Product datasheet for **MC209926**

### Batf (NM\_016767) Mouse Untagged Clone

#### Product data:

Product Type:	Expression Plasmids
Product Name:	Batf (NM_016767) Mouse Untagged Clone
Tag:	Tag Free
Symbol:	Batf
Synonyms:	B-ATF; SFA-2
Mammalian Cell Selection:	Neomycin
Vector:	pCMV6-Entry (PS100001)
E. coli Selection:	Kanamycin (25 ug/mL)
Fully Sequenced ORF:	>MC209926 representing NM_016767 Red=Cloning site Blue=ORF

TTTTGTAATACGACTCACTATAGGGCGGCCGGAATTCGTCGACTGGATCCGGTACCGAGGAGATCTGCC  
GCC**CGATCGCC**

ATGCCTCACAGCTCCGACAGCAGTGACTCCAGCTTCAGCCGCTCTCTCCCCCTGGCAAACAGGACTCAT  
CTGATGATGTGAGGAAAGTTCAGAGGAGAGAGAAGAATCGCATCGCTGCCAGAAGAGCCGACAGAGACA  
GACACAGAAAGCCGACACCCTTCACCTGGAGAGTGAGGACCTGGAGAAACAGAACGCAGCTCTCCGCAA  
GAGATCAAACAGCTCACCGAGGAGCTCAAGTACTTCACATCAGTGCTGAGCAGCCACGAGCCCTGTGCT  
CCGTGCTGGCCAGTGGCACCCCCTCGCCCCCGAGGTGGTATACAGTGCCCATGCCTTCCACCAGCCTCA  
CATCAGCTCGCCACGCTTCCAGCCCTGA

**ACGCGT**ACGCGGCCGCTCGAGCAGAAACTCATCTCAGAAGAGGATCTGGCAGCAAATGATATCCTGGATT  
ACAAGGATGACGACGATAAGGTTTAA

Restriction Sites:	Sgfl-MluI
ACCN:	NM_016767
Insert Size:	378 bp
OTI Disclaimer:	Our molecular clone sequence data has been matched to the reference identifier above as a point of reference. Note that the complete sequence of our molecular clones may differ from the sequence published for this corresponding reference, e.g., by representing an alternative RNA splicing form or single nucleotide polymorphism (SNP).



[View online »](#)

<b>Components:</b>	The ORF clone is ion-exchange column purified and shipped in a 2D barcoded Matrix tube containing 10ug of transfection-ready, dried plasmid DNA (reconstitute with 100 ul of water).
<b>Reconstitution Method:</b>	<ol style="list-style-type: none"><li>1. Centrifuge at 5,000xg for 5min.</li><li>2. Carefully open the tube and add 100ul of sterile water to dissolve the DNA.</li><li>3. Close the tube and incubate for 10 minutes at room temperature.</li><li>4. Briefly vortex the tube and then do a quick spin (less than 5000xg) to concentrate the liquid at the bottom.</li><li>5. Store the suspended plasmid at -20°C. The DNA is stable for at least one year from date of shipping when stored at -20°C.</li></ol>
<b>RefSeq:</b>	<a href="#">BC132410</a> , <a href="#">AAI32411</a>
<b>RefSeq Size:</b>	710 bp
<b>RefSeq ORF:</b>	378 bp
<b>Locus ID:</b>	53314
<b>UniProt ID:</b>	<a href="#">O35284</a>
<b>Cytogenetics:</b>	12 D2
<b>Gene Summary:</b>	AP-1 family transcription factor that controls the differentiation of lineage-specific cells in the immune system: specifically mediates the differentiation of T-helper 17 cells (Th17), follicular T-helper cells (TfH), CD8(+) dendritic cells and class-switch recombination (CSR) in B-cells. Acts via the formation of a heterodimer with JUNB that recognizes and binds DNA sequence 5'-TGA[CG]TCA-3'. The BATF-JUNB heterodimer also forms a complex with IRF4 (or IRF8) in immune cells, leading to recognition of AICE sequence (5'-TGAnTCA/GAAA-3'), an immune-specific regulatory element, followed by cooperative binding of BATF and IRF4 (or IRF8) and activation of genes. Controls differentiation of T-helper cells producing interleukin-17 (Th17 cells) by binding to Th17-associated gene promoters: regulates expression of the transcription factor RORC itself and RORC target genes such as IL17 (IL17A or IL17B). Also involved in differentiation of follicular T-helper cells (TfH) by directing expression of BCL6 and MAF. In B-cells, involved in class-switch recombination (CSR) by controlling the expression of both AICDA and of germline transcripts of the intervening heavy-chain region and constant heavy-chain region (I(H)-C(H)). Following infection, can participate in CD8(+) dendritic cell differentiation via interaction with IRF4 and IRF8 to mediate cooperative gene activation. Regulates effector CD8(+) T-cell differentiation by regulating expression of SIRT1. Following DNA damage, part of a differentiation checkpoint that limits self-renewal of hematopoietic stem cells (HSCs): up-regulated by STAT3, leading to differentiation of HSCs, thereby restricting self-renewal of HSCs.[UniProtKB/Swiss-Prot Function]