

Product datasheet for MC209979

Il1f6 (NM_019450) Mouse Untagged Clone

Product data:

Product Type:	Expression Plasmids
Product Name:	Il1f6 (NM_019450) Mouse Untagged Clone
Tag:	Tag Free
Symbol:	Il1f6
Synonyms:	Fil1; IL-1H1; Il1f9; IL1RP2; Il36a
Mammalian Cell Selection:	Neomycin
Vector:	pCMV6-Entry (PS100001)
E. coli Selection:	Kanamycin (25 ug/mL)
Fully Sequenced ORF:	>MC209979 representing NM_019450 Red=Cloning site Blue=ORF Orange=Stop codon

TTTTGTATACGACTCACTATAGGGCGGCCGGAATTCGTCGACTGGATCCGGTACCGAGGAGATCTGCC
 GCCCGCATCGCC

ATGAATAAGGAGAAAGAACTAAGAGCAGCATCACCTTCGCTTAGACATGTTCAGGATCTTAGTAGTCGTG
 TGTGGATCCTGCAGAACATATCCTCACTGCAGTCCCAAGGAAAGAGCAACAGTTCCAGTCACTATTAC
 CTTGCTCCCATGCCAATATCTGGACACTCTTGAGACGAACAGGGGGATCCACGTACATGGGAGTGCAA
 AGGCCGATGAGCTGCCTGTTCTGCACAAAGGATGGGAGCAGCCTGTGCTACAGCTTGGGAAGGGAACA
 TAATGGAAATGTACAACAAAAGGAACCTGTAAAAGCCTCTCTCTATCACAAGAAGAGTGGTACAAC
 CTCTACATTTGAGTCTGCAGCCTTCCCTGGTTGGTTCATCGCTGTCTGCTCTAAAGGGAGCTGCCCACTC
 ATTCTGACCCAAGAACTGGGGAAATCTTCATCACTGACTTCGAGATGATTGTGGTACATTAA

ACGCGTACGCGGCCGCTCGAGCAGAACTCATCTCAGAAGAGGATCTGGCAGCAAATGATATCCTGGATT
 ACAAGGATGACGACGATAAGGTTTAA

Restriction Sites:	Sgfl-MluI
ACCN:	NM_019450
Insert Size:	483 bp

OTI Disclaimer: Our molecular clone sequence data has been matched to the reference identifier above as a point of reference. Note that the complete sequence of our molecular clones may differ from the sequence published for this corresponding reference, e.g., by representing an alternative RNA splicing form or single nucleotide polymorphism (SNP).


[View online »](#)

Components:	The ORF clone is ion-exchange column purified and shipped in a 2D barcoded Matrix tube containing 10ug of transfection-ready, dried plasmid DNA (reconstitute with 100 ul of water).
Reconstitution Method:	<ol style="list-style-type: none"> 1. Centrifuge at 5,000xg for 5min. 2. Carefully open the tube and add 100ul of sterile water to dissolve the DNA. 3. Close the tube and incubate for 10 minutes at room temperature. 4. Briefly vortex the tube and then do a quick spin (less than 5000xg) to concentrate the liquid at the bottom. 5. Store the suspended plasmid at -20°C. The DNA is stable for at least one year from date of shipping when stored at -20°C.
Note:	Plasmids are not sterile. For experiments where strict sterility is required, filtration with 0.22um filter is required.
RefSeq:	<u>NM_019450.3</u> , <u>NP_062323.1</u>
RefSeq Size:	883 bp
RefSeq ORF:	483 bp
Locus ID:	54448
UniProt ID:	<u>Q9JLA2</u>
Cytogenetics:	2 16.26 cM
Gene Summary:	<p>Cytokine that binds to and signals through the IL1RL2/IL-36R receptor which in turn activates NF-kappa-B and MAPK signaling pathways in target cells linked to a pro-inflammatory response. Part of the IL-36 signaling system that is thought to be present in epithelial barriers and to take part in local inflammatory response; similar to the IL-1 system with which it shares the coreceptor IL1RAP. Seems to be involved in skin inflammatory response by acting on keratinocytes, dendritic cells and indirectly on T-cells to drive tissue infiltration, cell maturation and cell proliferation. Induces the production of proinflammatory cytokines, including IL-12, IL-1 beta, IL-6, TNF-alpha and IL-23 in bone marrow-derived dendritic cells (BMDCs). Involved in dendritic cell maturation by stimulating the surface expression of CD80, CD86 and MHC class II. Induces the production of IFN-gamma, IL-4 and IL-17 by cultured CD4(+) T-cells and splenocytes. May play a role in proinflammatory effects in the lung: induces the expression of CXCL1 and CXCL2 in the lung, and the expression of TNF-alpha, IL-36c, IL-1A, IL-1B, CXCL1 and CXCL2 in isolated splenic CD11c(+) alveolar macrophages. May be involved in T-cell maturation by stimulating the surface expression of CD40 and modestly CD80 and CD86 in splenic CD11c(+) cells. May be involved in CD4(+) T-cell proliferation. Induces NF-kappa B activation in macrophages.[UniProtKB/Swiss-Prot Function]</p>