

Product datasheet for MC210041

Hsd17b12 (NM_019657) Mouse Untagged Clone

Product data:

Product Type:	Expression Plasmids
Product Name:	Hsd17b12 (NM_019657) Mouse Untagged Clone
Tag:	Tag Free
Symbol:	Hsd17b12
Synonyms:	2610510O05Rik; AI172963; KIK-I; Kik1
Mammalian Cell Selection:	Neomycin
Vector:	pCMV6-Entry (PS100001)
E. coli Selection:	Kanamycin (25 ug/mL)
Fully Sequenced ORF:	>MC210041 representing NM_019657 Red=Cloning site Blue=ORF Orange=Stop codon

TTTGTAAATACGACTCACTATAGGGCGGCCGGAATTCGTCGACTGGATCCGGTACCGAGGAGATCTGCC
 GCC**CGATCGCC**

ATGGAGTGCCTCCCCGGCGCCGGCTTCTGTACTGGGTGGGCGGAGCACCATTGCCTACCTGGCCC
 TGCGAGCCTCGTACTCGCTCTTCAGGGCCTCCAGGTGTGGTGCCTGGGCAACGAGGCCTTGGTCGGTCC
 GCGACTCGGAGAATGGCAGTTGTTACAGGTGGCACTGATGGAATTGGAAAAGCATATGCAGAAGAGTTA
 GCAAAACGTGGAATGAAGATTGCTGATCAGCAGGTCCCAAGATAAACTGAATCAGTTTCCAACAACA
 TCAAAGAAAAATTCAACGTTGAAACAAGGACCATTGCTGTCGACTTTTCATTGGATGATATTTATGATAA
 AATTAAGACAGGCCTGTCTGGTCTTGAGATTGGCGTTTTAGTGAACAACGTGGGCATGTCATACGAGTAT
 CCGGAATACTTTCTGAAATTCCCGACTTGGACAACACCATCAAGAACTGATAAATATTAATGTGCTTT
 CCGTTTGCAAGGTGACACGCTTGGTGTGCTGGCATGGTAGAAAGATCTAAAGGGGTGATTCTCAACAT
 CTCCTCAGCCAGTGGCATGCTCCAGTTCATTGTTGACAATCTACTCTGCAACCAAGGCCTTTGTAGAT
 TTCTTCTCTCAGTGCCTCCATGAGGAGTATAAGAGCAAGGCATCTTTGTGCAGAGTGTGATGCCATACC
 TTGTAGCTACAACTGGCAAAAATACAGAAGCCGACTTTGGATAAGCCCTCTGCAGAGACATTTGTGAA
 GTCTGCAATTAACAGTAGGTTTGCAGACCCGAACCACTGGATATGTGATCCACTCTCATGGGCTCA
 ATAACTCAATCATGCCTCGTTGGATGTATTTTAAATAATCATGGGTTTCAGCAAGTCTTTCGCGGAATC
 GCTACCTGAAGAAAAGGAAGAAGAACTAA

ACGCGTACGCGGCCGCTCGAGCAGAACTCATCTCAGAAGAGGATCTGGCAGCAATGATATCCTGGATT
 ACAAGGATGACGACGATAAGGTTTAA

Restriction Sites:	SgfI-MluI
ACCN:	NM_019657


[View online »](#)

Insert Size:	939 bp
OTI Disclaimer:	<p>Due to the inherent nature of this plasmid, standard methods to replicate additional amounts of DNA in E. coli are highly likely to result in mutations and/or rearrangements. Therefore, OriGene does not guarantee the capability to replicate this plasmid DNA. Additional amounts of DNA can be purchased from OriGene with batch-specific, full-sequence verification at a reduced cost. Please contact our customer care team at custsupport@origene.com or by calling 301.340.3188 option 3 for pricing and delivery.</p> <p>The molecular sequence of this clone aligns with the gene accession number as a point of reference only. However, individual transcript sequences of the same gene can differ through naturally occurring variations (e.g. polymorphisms), each with its own valid existence. This clone is substantially in agreement with the reference, but a complete review of all prevailing variants is recommended prior to use. More info</p>
Components:	The ORF clone is ion-exchange column purified and shipped in a 2D barcoded Matrix tube containing 10ug of transfection-ready, dried plasmid DNA (reconstitute with 100 ul of water).
Reconstitution Method:	<ol style="list-style-type: none"> 1. Centrifuge at 5,000xg for 5min. 2. Carefully open the tube and add 100ul of sterile water to dissolve the DNA. 3. Close the tube and incubate for 10 minutes at room temperature. 4. Briefly vortex the tube and then do a quick spin (less than 5000xg) to concentrate the liquid at the bottom. 5. Store the suspended plasmid at -20°C. The DNA is stable for at least one year from date of shipping when stored at -20°C.
Note:	Plasmids are not sterile. For experiments where strict sterility is required, filtration with 0.22um filter is required.
RefSeq:	NM_019657.4 , NP_062631.1
RefSeq Size:	1839 bp
RefSeq ORF:	939 bp
Locus ID:	56348
UniProt ID:	O70503
Cytogenetics:	2 E1
Gene Summary:	<p>Catalyzes the second of the four reactions of the long-chain fatty acids elongation cycle. This endoplasmic reticulum-bound enzymatic process, allows the addition of two carbons to the chain of long- and very long-chain fatty acids/VLCFAs per cycle. This enzyme has a 3-ketoacyl-CoA reductase activity, reducing 3-ketoacyl-CoA to 3-hydroxyacyl-CoA, within each cycle of fatty acid elongation. Thereby, it may participate in the production of VLCFAs of different chain lengths that are involved in multiple biological processes as precursors of membrane lipids and lipid mediators. May also catalyze the transformation of estrone (E1) into estradiol (E2) and play a role in estrogen formation.[UniProtKB/Swiss-Prot Function]</p>