

## **Product datasheet for MC210277**

## Sarnp (NM\_025364) Mouse Untagged Clone

## **Product data:**

**Product Type:** Expression Plasmids

**Product Name:** Sarnp (NM\_025364) Mouse Untagged Clone

Tag: Tag Free
Symbol: Sarnp

**Synonyms:** 1110005A23Rik; Cip29

Mammalian Cell Neomycin

Selection:

Vector:

pCMV6-Entry (PS100001)

E. coli Selection: Kanamycin (25 ug/mL)

Fully Sequenced ORF: >MC210277 representing NM\_025364

Red=Cloning site Blue=ORF Orange=Stop codon

TTTTGTAATACGACTCACTATAGGGCGGCCGGGAATTCGTCGACTGGATCCGGTACCGAGGAGATCTGCC

GCCGCGATCGCC

TAA

ACGCGTACGCGGCCGCTCGAGCAGAAACTCATCTCAGAAGAGGATCTGGCAGCAAATGATATCCTGGATT

ACAAGGATGACGACGATAAGGTTTAA

**Restriction Sites:** Sgfl-Mlul ACCN: NM 025364

**Insert Size:** 633 bp



**OriGene Technologies, Inc.** 9620 Medical Center Drive, Ste 200

CN: techsupport@origene.cn

Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com



**OTI Disclaimer:** 

Our molecular clone sequence data has been matched to the reference identifier above as a point of reference. Note that the complete sequence of our molecular clones may differ from the sequence published for this corresponding reference, e.g., by representing an alternative RNA splicing form or single nucleotide polymorphism (SNP).

Components:

The ORF clone is ion-exchange column purified and shipped in a 2D barcoded Matrix tube containing 10ug of transfection-ready, dried plasmid DNA (reconstitute with 100 ul of water).

**Reconstitution Method:** 

- 1. Centrifuge at 5,000xg for 5min.
- 2. Carefully open the tube and add 100ul of sterile water to dissolve the DNA.
- 3. Close the tube and incubate for 10 minutes at room temperature.
- 4. Briefly vortex the tube and then do a quick spin (less than 5000xg) to concentrate the liquid

at the bottom.

5. Store the suspended plasmid at -20°C. The DNA is stable for at least one year from date of

shipping when stored at -20°C.

**Note:** Plasmids are not sterile. For experiments where strict sterility is required, filtration with

0.22um filter is required.

RefSeq: <u>NM 025364.2</u>, <u>NP 079640.1</u>

RefSeq Size: 1064 bp
RefSeq ORF: 633 bp
Locus ID: 66118
UniProt ID: Q9D1J3
Cytogenetics: 10 D3

**Gene Summary:** Binds both single-stranded and double-stranded DNA with higher affinity for the single-

stranded form. Specifically binds to scaffold/matrix attachment region DNA. Also binds single-stranded RNA. Enhances RNA unwinding activity of DDX39A. May participate in important transcriptional or translational control of cell growth, metabolism and carcinogenesis. Component of the TREX complex which is thought to couple mRNA transcription, processing and nuclear export, and specifically associates with spliced mRNA and not with unspliced premRNA. TREX is recruited to spliced mRNAs by a transcription-independent mechanism, binds to mRNA upstream of the exon-junction complex (EJC) and is recruited in a splicing- and capdependent manner to a region near the 5' end of the mRNA where it functions in mRNA export to the cytoplasm via the TAP/NFX1 pathway (By similarity).[UniProtKB/Swiss-Prot

Function]