

Product datasheet for MC210352

Lsm5 (NM 025520) Mouse Untagged Clone

Product data:

Product Type: Expression Plasmids

Product Name: Lsm5 (NM_025520) Mouse Untagged Clone

Tag: Tag Free
Symbol: Lsm5

Synonyms: 2010208O10Rik; 2310034K10Rik

Vector:pCMV6-Entry (PS100001)E. coli Selection:Kanamycin (25 ug/mL)

Cell Selection: Neomycin

Fully Sequenced ORF: >MC210352 representing NM_025520

Red=Cloning site Blue=ORF Orange=Stop codon

TTTTGTAATACGACTCACTATAGGGCGGCCGGGAATTCGTCGACTGGATCCGGTACCGAGGAGATCTGCC

GCCGCGATCGCC

ACGCGTACGCGGCCGCTCGAGCAGAAACTCATCTCAGAAGAGGATCTGGCAGCAAATGATATCCTGGATT

ACAAGGATGACGACGATAAGGTTTAA

Restriction Sites: Sgfl-Mlul

ACCN: NM_025520

Insert Size: 276 bp

OTI Disclaimer: Our molecular clone sequence data has been matched to the reference identifier above as a

point of reference. Note that the complete sequence of our molecular clones may differ from the sequence published for this corresponding reference, e.g., by representing an alternative

RNA splicing form or single nucleotide polymorphism (SNP).

Components: The ORF clone is ion-exchange column purified and shipped in a 2D barcoded Matrix tube

containing 10ug of transfection-ready, dried plasmid DNA (reconstitute with 100 ul of water).



OriGene Technologies, Inc. 9620 Medical Center Drive, Ste 200

CN: techsupport@origene.cn

Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com



Reconstitution Method:

- 1. Centrifuge at 5,000xg for 5min.
- 2. Carefully open the tube and add 100ul of sterile water to dissolve the DNA.
- 3. Close the tube and incubate for 10 minutes at room temperature.
- 4. Briefly vortex the tube and then do a quick spin (less than 5000xg) to concentrate the liquid

at the bottom.

5. Store the suspended plasmid at -20°C. The DNA is stable for at least one year from date of

shipping when stored at -20°C.

RefSeq: <u>NM 025520.3</u>, <u>NP 079796.1</u>

RefSeq Size: 516 bp
RefSeq ORF: 276 bp
Locus ID: 66373
UniProt ID: P62322
Cytogenetics: 6 B3

Gene Summary: Plays role in pre-mRNA splicing as component of the U4/U6-U5 tri-snRNP complex that is

involved in spliceosome assembly, and as component of the precatalytic spliceosome (spliceosome B complex). The heptameric LSM2-8 complex binds specifically to the 3'-

terminal U-tract of U6 snRNA.[UniProtKB/Swiss-Prot Function]