

## Product datasheet for MC210379

### Polr2l (NM\_025593) Mouse Untagged Clone

#### Product data:

Product Type:	Expression Plasmids
Product Name:	Polr2l (NM_025593) Mouse Untagged Clone
Tag:	Tag Free
Symbol:	Polr2l
Synonyms:	2510029B14Rik
Mammalian Cell Selection:	Neomycin
Vector:	pCMV6-Entry (PS100001)
E. coli Selection:	Kanamycin (25 ug/mL)
Fully Sequenced ORF:	>MC210379 representing NM_025593 Red=Cloning site Blue=ORF Orange=Stop codon

TTTGTAAATACGACTCACTATAGGGCGGCCGGAATTCGTCGACTGGATCCGGTACCGAGGAGATCTGCC  
GCC**CGATCGC**

ATGATCATCCCGGTGCGCTGCTTACCTGCGGGAAGATCGTCGGCAACAAATGGGAAGCCTACCTGGGTC  
TGCTGCAGGCCGAGTACACGGAGGGGGATGCCCTGGACGCTCTGGGCCTAAAGCGCTACTGCTGCCGCCG  
CATGCTGCTAGCACACGTGGACCTGATTGAGAAGCTGCTGAACATGCACCCCTAGAGAAG**TGA**

**ACGCGT**ACGCGGCCGCTCGAGCAGAACTCATCTCAGAAGAGGATCTGGCAGCAAATGATATCCTGGATT  
ACAAGGATGACGACGATAAGGTTTAA

Restriction Sites:	SgfI-MluI
ACCN:	NM_025593
Insert Size:	204 bp
OTI Disclaimer:	Our molecular clone sequence data has been matched to the reference identifier above as a point of reference. Note that the complete sequence of our molecular clones may differ from the sequence published for this corresponding reference, e.g., by representing an alternative RNA splicing form or single nucleotide polymorphism (SNP).
Components:	The ORF clone is ion-exchange column purified and shipped in a 2D barcoded Matrix tube containing 10ug of transfection-ready, dried plasmid DNA (reconstitute with 100 ul of water).


[View online »](#)

**Reconstitution Method:**

1. Centrifuge at 5,000xg for 5min.
2. Carefully open the tube and add 100ul of sterile water to dissolve the DNA.
3. Close the tube and incubate for 10 minutes at room temperature.
4. Briefly vortex the tube and then do a quick spin (less than 5000xg) to concentrate the liquid at the bottom.
5. Store the suspended plasmid at -20°C. The DNA is stable for at least one year from date of shipping when stored at -20°C.

**RefSeq:** [NM\\_025593.1](#), [NP\\_079869.1](#)

**RefSeq Size:** 1671 bp

**RefSeq ORF:** 204 bp

**Locus ID:** 66491

**UniProt ID:** [P62876](#)

**Cytogenetics:** 7 F5

**Gene Summary:** DNA-dependent RNA polymerase catalyzes the transcription of DNA into RNA using the four ribonucleoside triphosphates as substrates. Common component of RNA polymerases I, II and III which synthesize ribosomal RNA precursors, mRNA precursors and many functional non-coding RNAs, and a small RNAs, such as 5S rRNA and tRNAs, respectively. Pol II is the central component of the basal RNA polymerase II transcription machinery. Pols are composed of mobile elements that move relative to each other. In Pol II, POLR2L/RBP10 is part of the core element with the central large cleft (By similarity).[UniProtKB/Swiss-Prot Function]