

Product datasheet for MC210383

Chtop (NM_023215) Mouse Untagged Clone

Product data:

Product Type: Expression Plasmids

Product Name: Chtop (NM_023215) Mouse Untagged Clone

Tag: Tag Free Symbol: Chtop

Synonyms: 2500003M10Rik; Fop; Srag

Mammalian Cell Neomycin

Selection:

Vector:pCMV6-Entry (PS100001)E. coli Selection:Kanamycin (25 ug/mL)

Fully Sequenced ORF: >MC210383 representing NM_023215

Red=Cloning site Blue=ORF Orange=Stop codon

TTTTGTAATACGACTCACTATAGGGCGGCCGGGAATTCGTCGACTGGATCCGGTACCGAGGAGATCTGCC

GCCGCGATCGCC

TTGGATGCCTACATGGCACAGACAGATCCTGAAACCAATGAT<mark>TGA</mark>

 ACGCGTACGCGGCCGCTCGAGCAGAAACTCATCTCAGAAGAGGATCTGGCAGCAAATGATATCCTGGATT

ACAAGGATGACGACGATAAGGTTTAA

Restriction Sites: Sgfl-Mlul

ACCN: NM_023215

Insert Size: 675 bp



OriGene Technologies, Inc. 9620 Medical Center Drive, Ste 200

CN: techsupport@origene.cn

Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com ORIGENE

OTI Disclaimer:

Our molecular clone sequence data has been matched to the reference identifier above as a point of reference. Note that the complete sequence of our molecular clones may differ from the sequence published for this corresponding reference, e.g., by representing an alternative RNA splicing form or single nucleotide polymorphism (SNP).

Components:

The ORF clone is ion-exchange column purified and shipped in a 2D barcoded Matrix tube containing 10ug of transfection-ready, dried plasmid DNA (reconstitute with 100 ul of water).

Reconstitution Method:

- 1. Centrifuge at 5,000xg for 5min.
- 2. Carefully open the tube and add 100ul of sterile water to dissolve the DNA.
- 3. Close the tube and incubate for 10 minutes at room temperature.
- 4. Briefly vortex the tube and then do a quick spin (less than 5000xg) to concentrate the liquid

at the bottom.

5. Store the suspended plasmid at -20°C. The DNA is stable for at least one year from date of

shipping when stored at -20°C.

Note: Plasmids are not sterile. For experiments where strict sterility is required, filtration with

0.22um filter is required.

RefSeq: <u>NM 023215.6</u>, <u>NP 075704.2</u>

3 F1

 RefSeq Size:
 3721 bp

 RefSeq ORF:
 675 bp

 Locus ID:
 66511

 UniProt ID:
 Q9CY57

Gene Summary:

Cytogenetics:

Plays an important role in the ligand-dependent activation of estrogen receptor target genes (By similarity). May play a role in the silencing of fetal globin genes (PubMed:20688955). Recruits the 5FMC complex to ZNF148, leading to desumoylation of ZNF148 and subsequent transactivation of ZNF148 target genes (PubMed:22872859). Required for the tumorigenicity of glioblastoma cells. Binds to 5-hydroxymethylcytosine (5hmC) and associates with the methylosome complex containing PRMT1, PRMT5, MEP50 and ERH. The CHTOP-methylosome complex associated with 5hmC methylates H4R3 and transactivates genes involved in glioblastomagenesis (PubMed:25284789).[UniProtKB/Swiss-Prot Function]

Transcript Variant: This variant (3) differs in the 5' UTR, lacks a portion of the 5' coding region, and uses a downstream start codon compared to variant 1. The encoded isoform (3) has a shorter N-terminus compared to isoform 1. Variants 3 and 8 encode the same isoform. Sequence Note: The RefSeq transcript and protein were derived from genomic sequence to make the sequence consistent with the reference genome assembly. The genomic

coordinates used for the transcript record were based on alignments.