

## Product datasheet for MC210410

### Tsen15 (NM\_025677) Mouse Untagged Clone

#### Product data:

Product Type: Expression Plasmids

Product Name: Tsen15 (NM\_025677) Mouse Untagged Clone

Tag: Tag Free

Symbol: Tsen15

Synonyms: 5730449L18Rik; AL023077; Sen15

Mammalian Cell Selection: Neomycin

Vector: pCMV6-Entry (PS100001)

E. coli Selection: Kanamycin (25 ug/mL)

Fully Sequenced ORF: >MC210410 representing NM\_025677  
Red=Cloning site Blue=ORF Orange=Stop codon

TTTGTAAATACGACTCACTATAGGGCGGCCGGAATTCGTCGACTGGATCCGGTACCGAGGAGATCTGCC  
GCC**CGATCGCC**

ATGGAGGAGCGCAGCGATTCCAAACCTACCCGGGATGTAGCGGGCCTGGCCGGCTCCCGTTTCGCGATG  
GCGGCGGCGCCACACATGGGCTCCGGAGGACGCCTGGATGGGCACACACCCTAAGTATTTAGAAATGAT  
GGAATTAGATATAGGAGATGCCACCAAGTTTATATAGCATTCTTGTTTACCTGGATCTCATGGAGAGT  
AAAAGTTGGCATGAAGTAACTGTGTAGGAATACCAGAACTACAACCTCATCTGCCTCCTTGGCACTGAGA  
TCGAAGGGGAAGGGCTGCAGACGGTGGTGCCTACACCCATTTCTGCTTCCCTCAGCCATAATAGGATAAG  
GGAAATCTTGAAGGCGTCTAGAAAGTTGAAGGCATCCAGAACTGCCGATGTCTTTACTTTGGCCATA  
GTGGAGTCAGATTCCACAATAGTCTATTATAAACTTACCGATGGATTTATGCTGCCAGACCTCAGAATA  
TTTCTCTTAGAAG**TGA**

**ACGCGT**ACGCGGCCGCTCGAGCAGAACTCATCTCAGAAGAGGATCTGGCAGCAAATGATATCCTGGATT  
ACAAGGATGACGACGATAAGGTTTAA

Restriction Sites: SgfI-MluI

ACCN: NM\_025677

Insert Size: 507 bp

OTI Disclaimer: Our molecular clone sequence data has been matched to the reference identifier above as a point of reference. Note that the complete sequence of our molecular clones may differ from the sequence published for this corresponding reference, e.g., by representing an alternative RNA splicing form or single nucleotide polymorphism (SNP).



[View online »](#)

<b>Components:</b>	The ORF clone is ion-exchange column purified and shipped in a 2D barcoded Matrix tube containing 10ug of transfection-ready, dried plasmid DNA (reconstitute with 100 ul of water).
<b>Reconstitution Method:</b>	<ol style="list-style-type: none"> <li>1. Centrifuge at 5,000xg for 5min.</li> <li>2. Carefully open the tube and add 100ul of sterile water to dissolve the DNA.</li> <li>3. Close the tube and incubate for 10 minutes at room temperature.</li> <li>4. Briefly vortex the tube and then do a quick spin (less than 5000xg) to concentrate the liquid at the bottom.</li> <li>5. Store the suspended plasmid at -20°C. The DNA is stable for at least one year from date of shipping when stored at -20°C.</li> </ol>
<b>Note:</b>	Plasmids are not sterile. For experiments where strict sterility is required, filtration with 0.22um filter is required.
<b>RefSeq:</b>	<a href="#">NM_025677.3</a> , <a href="#">NP_079953.2</a>
<b>RefSeq Size:</b>	1147 bp
<b>RefSeq ORF:</b>	507 bp
<b>Locus ID:</b>	66637
<b>UniProt ID:</b>	<a href="#">Q8R3W5</a>
<b>Cytogenetics:</b>	1 G2
<b>Gene Summary:</b>	Non-catalytic subunit of the tRNA-splicing endonuclease complex, a complex responsible for identification and cleavage of the splice sites in pre-tRNA. It cleaves pre-tRNA at the 5' and 3' splice sites to release the intron. The products are an intron and two tRNA half-molecules bearing 2',3' cyclic phosphate and 5'-OH termini. There are no conserved sequences at the splice sites, but the intron is invariably located at the same site in the gene, placing the splice sites an invariant distance from the constant structural features of the tRNA body. The tRNA splicing endonuclease is also involved in mRNA processing via its association with pre-mRNA 3'-end processing factors, establishing a link between pre-tRNA splicing and pre-mRNA 3'-end formation, suggesting that the endonuclease subunits function in multiple RNA-processing events (By similarity).[UniProtKB/Swiss-Prot Function]