

Product datasheet for MC210546

Pdlim7 (NM_001114087) Mouse Untagged Clone

Product data:

Product Type: Expression Plasmids

Product Name: Pdlim7 (NM_001114087) Mouse Untagged Clone

Tag: Tag Free
Symbol: Pdlim7
Synonyms: LMP

Vector:pCMV6-Entry (PS100001)E. coli Selection:Kanamycin (25 ug/mL)

Cell Selection: Neomycin

Fully Sequenced ORF: >MC210546 representing NM_001114087

Red=Cloning site Blue=ORF Orange=Stop codon

TTTTGTAATACGACTCACTATAGGGCGGCCGGGAATTCGTCGACTGGATCCGGTACCGAGGAGATCTGCC

GCCGCGATCGCC

ACGCGTACGCGGCCGCTCGAGCAGAAACTCATCTCAGAAGAGGATCTGGCAGCAAATGATATCCTGGATT

ACAAGGATGACGACGATAAGGTTTAA

Restriction Sites: Sgfl-Mlul

ACCN: NM_001114087

Insert Size: 669 bp



OriGene Technologies, Inc. 9620 Medical Center Drive, Ste 200

CN: techsupport@origene.cn

Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com

Pdlim7 (NM_001114087) Mouse Untagged Clone - MC210546

OTI Disclaimer: Our molecular clone sequence data has been matched to the reference identifier above as a

point of reference. Note that the complete sequence of our molecular clones may differ from the sequence published for this corresponding reference, e.g., by representing an alternative

RNA splicing form or single nucleotide polymorphism (SNP).

Components: The ORF clone is ion-exchange column purified and shipped in a 2D barcoded Matrix tube

containing 10ug of transfection-ready, dried plasmid DNA (reconstitute with 100 ul of water).

Reconstitution Method: 1. Centrifuge at 5,000xg for 5min.

2. Carefully open the tube and add 100ul of sterile water to dissolve the DNA.

3. Close the tube and incubate for 10 minutes at room temperature.

4. Briefly vortex the tube and then do a quick spin (less than 5000xg) to concentrate the liquid

at the bottom.

5. Store the suspended plasmid at -20°C. The DNA is stable for at least one year from date of

shipping when stored at -20°C.

RefSeq: NM 001114087.2, NP 001107559.1

RefSeq Size: 1014 bp
RefSeq ORF: 669 bp
Locus ID: 67399
Cytogenetics: 13 B1

Gene Summary: May function as a scaffold on which the coordinated assembly of proteins can occur. May

play a role as an adapter that, via its PDZ domain, localizes LIM-binding proteins to actin filaments of both skeletal muscle and nonmuscle tissues. Involved in both of the two fundamental mechanisms of bone formation, direct bone formation (e.g. embryonic flat bones mandible and cranium), and endochondral bone formation (e.g. embryonic long bone development). Plays a role during fracture repair. Involved in BMP6 signaling pathway (By

similarity).[UniProtKB/Swiss-Prot Function]

Transcript Variant: This variant (b) lacks several exons, and uses an alternate 3'-terminal exon, compared to variant a. This results in a novel 3' coding region and 3' UTR, compared to

variant a. The encoded isoform (b) has a shorter and distinct C-terminus, compared to

isoform a.