

Product datasheet for MC210623

N6amt1 (NM_001159331) Mouse Untagged Clone

Product data:

Product Type: Expression Plasmids

Product Name: N6amt1 (NM_001159331) Mouse Untagged Clone

Tag: Tag Free Symbol: N6amt1

Synonyms: 5830445C04Rik; Hemk2; Pred28

Mammalian Cell

Selection:

Neomycin

Vector:pCMV6-Entry (PS100001)E. coli Selection:Kanamycin (25 ug/mL)

Fully Sequenced ORF: >MC210623 representing NM_001159331

Red=Cloning site Blue=ORF Orange=Stop codon

TTTTGTAATACGACTCACTATAGGGCGGCCGGGAATTCGTCGACTGGATCCGGTACCGAGGAGATCTGCC

GCCGCGATCGCC

ACGCGTACGCGGCCGCTCGAGCAGAAACTCATCTCAGAAGAGGATCTGGCAGCAAATGATATCCTGGATT

ACAAGGATGACGACGATAAGGTTTAA

Restriction Sites: Sgfl-Mlul

ACCN: NM_001159331

Insert Size: 417 bp

OTI Disclaimer: Our molecular clone sequence data has been matched to the reference identifier above as a

point of reference. Note that the complete sequence of our molecular clones may differ from the sequence published for this corresponding reference, e.g., by representing an alternative

RNA splicing form or single nucleotide polymorphism (SNP).



OriGene Technologies, Inc. 9620 Medical Center Drive, Ste 200

CN: techsupport@origene.cn

Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com



Components:

The ORF clone is ion-exchange column purified and shipped in a 2D barcoded Matrix tube containing 10ug of transfection-ready, dried plasmid DNA (reconstitute with 100 ul of water).

Reconstitution Method:

- 1. Centrifuge at 5,000xg for 5min.
- 2. Carefully open the tube and add 100ul of sterile water to dissolve the DNA.
- 3. Close the tube and incubate for 10 minutes at room temperature.
- 4. Briefly vortex the tube and then do a quick spin (less than 5000xg) to concentrate the liquid

at the bottom.

5. Store the suspended plasmid at -20°C. The DNA is stable for at least one year from date of

shipping when stored at -20°C.

Note: Plasmids are not sterile. For experiments where strict sterility is required, filtration with

0.22um filter is required.

RefSeq: NM 001159331.1, NP 001152803.1

RefSeq Size: 1649 bp RefSeq ORF: 417 bp Locus ID: 67768 **UniProt ID:** Q6SKR2 Cytogenetics: 16 C3.3

Gene Summary: Methyltransferase that can methylate both proteins and DNA, and to a lower extent, arsenic

> (PubMed:20606008, PubMed:26797129). Catalytic subunit of a heterodimer with TRMT112, which catalyzes N5-methylation of Glu residue of proteins with a Gly-Gln-Xaa-Xaa-Arg motif (PubMed:26797129). Methylates ETF1 on 'Gln-185'; ETF1 needs to be complexed to ERF3 in its GTP-bound form to be efficiently methylated (PubMed:20606008, PubMed:26797129). Also acts as a N(6)-adenine-specific DNA methyltransferase by mediating methylation of DNA on the 6th position of adenine (N(6)-methyladenosine) (By similarity). N(6)-methyladenosine

(m6A) DNA is significantly enriched in exonic regions and is associated with gene

transcriptional activation (By similarity). May also play a role in the modulation of arsenicinduced toxicity by mediating the conversion of monomethylarsonous acid (3+) into the less toxic dimethylarsonic acid (By similarity). It however only plays a limited role in arsenic

metabolism compared with AS3MT (By similarity).[UniProtKB/Swiss-Prot Function] Transcript Variant: This variant (2) lacks an alternate coding exon compared to variant 1, that causes a frameshift. The resulting isoform (2) has a shorter and distinct C-terminus compared

to isoform 1. Sequence Note: This RefSeq record was created from transcript and genomic sequence data to make the sequence consistent with the reference genome assembly. The genomic coordinates used for the transcript record were based on transcript alignments.