

## Product datasheet for MC210696

### Eif3h (NM\_080635) Mouse Untagged Clone

#### Product data:

Product Type:	Expression Plasmids
Product Name:	Eif3h (NM_080635) Mouse Untagged Clone
Tag:	Tag Free
Symbol:	Eif3h
Synonyms:	40kD; 1110008A16Rik; 9430017H16Rik; EIF3-gamma; EIF3-P40; Eif3s3
Mammalian Cell Selection:	Neomycin
Vector:	pCMV6-Entry (PS100001)
E. coli Selection:	Kanamycin (25 ug/mL)
Fully Sequenced ORF:	>MC210696 representing NM_080635 Red=Cloning site Blue=ORF Orange=Stop codon

TTTTGTAATACGACTCACTATAGGGCGGCCGGAATTCGTCGACTGGATCCGGTACCGAGGAGATCTGCC  
GCC**CGATCGCC**

ATGGCGTCGCGCAAGGAAGGCACCGGCTCTACCGCACCTCCTCCGGTTCTGCTGGCGGCGCGGTGGGA  
AGGGCAAAGGCAAAGGCGGCTCCGGAGATTCGGCCGTGAAGCAGGTGCAGATCGACGGCCTGGTAGTATT  
AAAGATAATCAAACATTATCAAGAAGAAGGACAAGGCACTGAGGTTGTTCCAGGGCGTCTCTGGGTCTG  
GTTGTGGAAGACCGGCTAGAGATTACCAACTGCTTCCCGTCCCCAGCACACGGAGGATGATGCTGACT  
TTGATGAGGTACAGTATCAGATGGAGATGATGCGCAGCCTTCGCCATGTCAACATTGATCACCTCCACGT  
GGGCTGGTATCAGTCCACATATTACGGCTCCTTGTTACCCGGGCGCTTCTGGATTCTCAGTTCAGCTAC  
CAGCACGCCATTGAAGAGTCTGTCTCCTCATTATGATCCCATAAAAACTGCCAAGGATCTCTCTCGC  
TGAAGGCGTACAGACTGACTCCTAAACTGATGGAAGTTTGTAAAGAGAAGGACTTTCCCTGAAGCATT  
GAAAAAGGCAAGCATCACCTTTGAGCACATGTTGAAGAAGTGCCGATTGTAATAAAACTCACATCTG  
ATCAATGTCCTTATGTGGGAAGTGAAGAAGTCAAGTCAAGTCAAGTCAAGTCAAGTCAAGTCAAGTCAAG  
CTAGCAGCAATCATCTGGGAAGAGCCTCCAGCTGCTGATGGACCGGTGGACGAAATGAGCCAGGACAT  
AATCAAATACAACACGTACATGCGCAACACCAGTAAGCAGCAGCAGCAGAAACATCAGTATCAGCAGCGT  
CGCCAACAGGAGAATATGCAGCGACAGAGTCGAGGCGAGCCCCACTCCCTGAGGAGGACTCTCTAAAC  
TCTTCAAGCCCCACCAGGCCCTGCCAGGATGGATTCACTGCTCATTGCAGGCCAGATTAACACTTACTG  
CCAGAACATCAAGGAGTTCAGTCCCAAACTTAGGCAAACCTTTCATGGCCAGGCTCTCAAGAATAC  
AATAATTAA

**ACGCGT**ACGCGGCCGCTCGAGCAGAACTCATCTCAGAAGAGGATCTGGCAGCAAATGATATCCTGGATT  
ACAAGGATGACGACGATAAGGTTTAA

Restriction Sites: SgfI-MluI



[View online >](#)

<b>ACCN:</b>	NM_080635
<b>Insert Size:</b>	1059 bp
<b>OTI Disclaimer:</b>	Our molecular clone sequence data has been matched to the reference identifier above as a point of reference. Note that the complete sequence of our molecular clones may differ from the sequence published for this corresponding reference, e.g., by representing an alternative RNA splicing form or single nucleotide polymorphism (SNP).
<b>Components:</b>	The ORF clone is ion-exchange column purified and shipped in a 2D barcoded Matrix tube containing 10ug of transfection-ready, dried plasmid DNA (reconstitute with 100 ul of water).
<b>Reconstitution Method:</b>	<ol style="list-style-type: none"><li>1. Centrifuge at 5,000xg for 5min.</li><li>2. Carefully open the tube and add 100ul of sterile water to dissolve the DNA.</li><li>3. Close the tube and incubate for 10 minutes at room temperature.</li><li>4. Briefly vortex the tube and then do a quick spin (less than 5000xg) to concentrate the liquid at the bottom.</li><li>5. Store the suspended plasmid at -20°C. The DNA is stable for at least one year from date of shipping when stored at -20°C.</li></ol>
<b>RefSeq:</b>	<a href="#">NM_080635.1</a> , <a href="#">NP_542366.1</a>
<b>RefSeq Size:</b>	1254 bp
<b>RefSeq ORF:</b>	1059 bp
<b>Locus ID:</b>	68135
<b>UniProt ID:</b>	<a href="#">Q91WK2</a>
<b>Cytogenetics:</b>	15 C
<b>Gene Summary:</b>	Component of the eukaryotic translation initiation factor 3 (eIF-3) complex, which is required for several steps in the initiation of protein synthesis. The eIF-3 complex associates with the 40S ribosome and facilitates the recruitment of eIF-1, eIF-1A, eIF-2:GTP:methionyl-tRNA <sup>i</sup> and eIF-5 to form the 43S pre-initiation complex (43S PIC). The eIF-3 complex stimulates mRNA recruitment to the 43S PIC and scanning of the mRNA for AUG recognition. The eIF-3 complex is also required for disassembly and recycling of post-termination ribosomal complexes and subsequently prevents premature joining of the 40S and 60S ribosomal subunits prior to initiation. The eIF-3 complex specifically targets and initiates translation of a subset of mRNAs involved in cell proliferation, including cell cycling, differentiation and apoptosis, and uses different modes of RNA stem-loop binding to exert either translational activation or repression.[UniProtKB/Swiss-Prot Function]