

Product datasheet for MC211349

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

OriGene Technologies, Inc.

2810006K23Rik (NM 028310) Mouse Untagged Clone

Product data:

Product Type: Expression Plasmids

Product Name: 2810006K23Rik (NM_028310) Mouse Untagged Clone

Tag: Tag Free

Symbol: 2810006K23Rik

Vector: pCMV6-Entry (PS100001) **E. coli Selection:** Kanamycin (25 ug/mL)

Cell Selection: Neomycin

Fully Sequenced ORF: >MC211349 representing NM_028310

Red=Cloning site Blue=ORF Orange=Stop codon

TTTTGTAATACGACTCACTATAGGGCGGCCGGGAATTCGTCGACTGGATCCGGTACCGAGGAGATCTGCC

GCCGCGATCGCC

ACGCGTACGCGGCCGCTCGAGCAGAAACTCATCTCAGAAGAGGATCTGGCAGCAAATGATATCCTGGATT

ACAAGGATGACGACGATAAGGTTTAA

Restriction Sites: Sgfl-Mlul
ACCN: NM_028310
Insert Size: 387 bp

OTI Disclaimer: Our molecular clone sequence data has been matched to the reference identifier above as a

point of reference. Note that the complete sequence of our molecular clones may differ from the sequence published for this corresponding reference, e.g., by representing an alternative

RNA splicing form or single nucleotide polymorphism (SNP).

Components: The ORF clone is ion-exchange column purified and shipped in a 2D barcoded Matrix tube

containing 10ug of transfection-ready, dried plasmid DNA (reconstitute with 100 ul of water).





Reconstitution Method:

- 1. Centrifuge at 5,000xg for 5min.
- 2. Carefully open the tube and add 100ul of sterile water to dissolve the DNA.
- 3. Close the tube and incubate for 10 minutes at room temperature.
- 4. Briefly vortex the tube and then do a quick spin (less than 5000xg) to concentrate the liquid at the bottom.
- 5. Store the suspended plasmid at -20°C. The DNA is stable for at least one year from date of shipping when stored at -20°C.

RefSeq: <u>NM 028310.3, NP 082586.1</u>

RefSeq Size: 1636 bp
RefSeq ORF: 387 bp
Locus ID: 72650
UniProt ID: Q80VP5
Cytogenetics: 5 F

Gene Summary: May act as a codon-independent translation release factor that has lost all stop codon

specificity and directs the termination of translation in mitochondrion. May help rescuing stalled mitoribosomes during translation (By similarity).[UniProtKB/Swiss-Prot Function] Transcript Variant: This variant (2) contains an alternate 3' terminal exon that extends past a splice site that is used in variant 1. This results in a novel 3' coding region and 3' UTR, compared to variant 1. It encodes isoform b which is shorter and has a distinct C-terminus,

compared to isoform a.