

Product datasheet for MC211471

Psmb11 (NM_175204) Mouse Untagged Clone

Product data:

Product Type: Expression Plasmids

Product Name: Psmb11 (NM_175204) Mouse Untagged Clone

Tag: Tag Free
Symbol: Psmb11

Synonyms: 5830406J20Rik; beta5t

Mammalian Cell Nec

Selection:

Neomycin

Vector:pCMV6-Entry (PS100001)E. coli Selection:Kanamycin (25 ug/mL)

Restriction Sites: Sgfl-Mlul ACCN: NM 175204

Insert Size: 909 bp

OTI Disclaimer: Our molecular clone sequence data has been matched to the reference identifier above as a

point of reference. Note that the complete sequence of our molecular clones may differ from the sequence published for this corresponding reference, e.g., by representing an alternative

RNA splicing form or single nucleotide polymorphism (SNP).

Components: The ORF clone is ion-exchange column purified and shipped in a 2D barcoded Matrix tube

containing 10ug of transfection-ready, dried plasmid DNA (reconstitute with 100 ul of water).

Reconstitution Method: 1. Centrifuge at 5,000xg for 5min.

2. Carefully open the tube and add 100ul of sterile water to dissolve the DNA.

3. Close the tube and incubate for 10 minutes at room temperature.

4. Briefly vortex the tube and then do a quick spin (less than 5000xg) to concentrate the liquid

at the bottom.

5. Store the suspended plasmid at -20°C. The DNA is stable for at least one year from date of

shipping when stored at -20°C.

RefSeq: NM 175204.4, NP 780413.1

RefSeq Size: 4247 bp RefSeq ORF: 909 bp



OriGene Technologies, Inc. 9620 Medical Center Drive, Ste 200

CN: techsupport@origene.cn

Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com



Psmb11 (NM_175204) Mouse Untagged Clone - MC211471

Locus ID: 73902

UniProt ID: Q8BG41

Cytogenetics: 14 C2

Gene Summary: The proteasome is a multicatalytic proteinase complex which is characterized by its ability to

cleave peptides with Arg, Phe, Tyr, Leu, and Glu adjacent to the leaving group at neutral or slightly basic pH. The proteasome has an ATP-dependent proteolytic activity. Incorporated

instead of PSMB5 or PSMB8, this unit reduces the chymotrypsin-like activity of the proteasome. Plays a pivotal role in development of CD8-positive T-cells.[UniProtKB/Swiss-

Prot Function]