

## Product datasheet for MC211780

### Calcoco2 (NM\_029755) Mouse Untagged Clone

#### Product data:

**Product Type:** Expression Plasmids  
**Product Name:** Calcoco2 (NM\_029755) Mouse Untagged Clone  
**Tag:** Tag Free  
**Symbol:** Calcoco2  
**Synonyms:** 2410154J16Rik; C77254; Ndp52; Ndp52l1  
**Mammalian Cell Selection:** Neomycin  
**Vector:** pCMV6-Entry (PS100001)  
**E. coli Selection:** Kanamycin (25 ug/mL)  
**Fully Sequenced ORF:** >MC211780 representing NM\_029755  
 Red=Cloning site Blue=ORF Orange=Stop codon

TTTTGTAATACGACTCACTATAGGGCGGCCGGAATTCGTCGACTGGATCCGGTACCGAGGAGATCTGCC  
 GCCCGCATCGCC

ATGGACCAGTGCCCCATACCTACCTTGCTGGAACATGGCAACTTCTCTCAGGTCCTGTTTAAACAATGTGG  
 AGAAGTTCTATGCTCCTAGAGGAGATATCATGTGCTATTACACCCTCACTGAAAAGTTTCATCCCTCGACG  
 CAAGGACTGGATTGGCATCTTTAAAGTAGGGTGAAGACCACTCAGGAGTATTATACCTTCATGTGGCT  
 CCCTTGCCAAAAGACAAAACAAGGATTCAGCCACACAGCAGGAAATCCAATTCAAAGCTTATTACCTC  
 CCAAGGATGTGGAGCGCTACCAGTCTGTATGTGGATGAAGATGGTTTAGTCCGGGGAACAAGTGTCCC  
 TTTCCAGTTTTGTCCAGACCCTGACGAGGACATAATGGTTGTTATCAATAAGGAAAAGGTAGAAGAGATG  
 GAACAGCTCAGTGAGGAGCTTTACCAACAAAACCAGGAGCTGAAAGACAAGTACGCTGACCTCCATGAGC  
 AGCTACAGAGGAAGCAGGTGGCACTGGAAGCAACAGAGGGTCAATAAGACCTTAGAACACAAAGTGA  
 AGAGAAGGCCCTCTGGGAGAAAGAGAAGGCCCTCTGGGAGGAAGAGAAGGCCCTCTGGGAGGAAGAGA  
 GCCTCTGGGAGGAAGAGAAGGCCCTCTGGGAGGAAGAGAAGGCCCTCTGGGAGGAAGAGAAGGCCCTCT  
 GGGAGGAAGAGAAGGCCCTCTGGGAGGAAGAGAAGGCCCTCTGGGAGGAAGAGAAGGCCCTCTGGGAGGA  
 AGAGAAGGCCCTCTGGGAGGAAGAGAAGGCCCTCTGGGAGAAAGAGAAGGCCCTCTGGGAGGAAGAGA  
 GCCTCTGGGAGAAAGAGAAGGCCCTGGGAGGTAGAGAAGGCCCTGGAAGGAAGTGAAGGCCATT  
 GGTGGAATGATCTGCACCGGTGA

ACGCGTACGCGGCCGCTCGAGCAGAAACTCATCTCAGAAGAGGATCTGGCAGCAAATGATATCTGGATT  
 ACAAGGATGACGACGATAAGGTTTAA

**Restriction Sites:** SgfI-MluI  
**ACCN:** NM\_029755



[View online »](#)

<b>Insert Size:</b>	933 bp
<b>OTI Disclaimer:</b>	Our molecular clone sequence data has been matched to the reference identifier above as a point of reference. Note that the complete sequence of our molecular clones may differ from the sequence published for this corresponding reference, e.g., by representing an alternative RNA splicing form or single nucleotide polymorphism (SNP).
<b>Components:</b>	The ORF clone is ion-exchange column purified and shipped in a 2D barcoded Matrix tube containing 10ug of transfection-ready, dried plasmid DNA (reconstitute with 100 ul of water).
<b>Reconstitution Method:</b>	<ol style="list-style-type: none"><li>1. Centrifuge at 5,000xg for 5min.</li><li>2. Carefully open the tube and add 100ul of sterile water to dissolve the DNA.</li><li>3. Close the tube and incubate for 10 minutes at room temperature.</li><li>4. Briefly vortex the tube and then do a quick spin (less than 5000xg) to concentrate the liquid at the bottom.</li><li>5. Store the suspended plasmid at -20°C. The DNA is stable for at least one year from date of shipping when stored at -20°C.</li></ol>
<b>RefSeq:</b>	<u><a href="#">NM_029755.3</a></u> , <u><a href="#">NP_084031.2</a></u>
<b>RefSeq Size:</b>	1556 bp
<b>RefSeq ORF:</b>	933 bp
<b>Locus ID:</b>	76815
<b>Cytogenetics:</b>	11 59.4 cM
<b>Gene Summary:</b>	Xenophagy-specific receptor required for autophagy-mediated intracellular bacteria degradation (By similarity). Acts as an effector protein of galectin-sensed membrane damage that restricts the proliferation of infecting pathogens upon entry into the cytosol by targeting LGALS8-associated bacteria for autophagy (By similarity). Initially orchestrates bacteria targeting to autophagosomes and subsequently ensures pathogen degradation by regulating pathogen-containing autophagosome maturation (By similarity). Bacteria targeting to autophagosomes relies on its interaction with MAP1LC3A, MAP1LC3B and/or GABARAPL2, whereas regulation of pathogen-containing autophagosome maturation requires the interaction with MAP3LC3C (By similarity). May play a role in ruffle formation and actin cytoskeleton organization and seems to negatively regulate constitutive secretion (By similarity).[UniProtKB/Swiss-Prot Function]