

Product datasheet for **MC212048**

Prmt6 (NM_178891) Mouse Untagged Clone

Product data:

Product Type: Expression Plasmids
Product Name: Prmt6 (NM_178891) Mouse Untagged Clone
Tag: Tag Free
Symbol: Prmt6
Synonyms: AW124876; BB233495; Hrmt116
Vector: pCMV6-Entry (PS100001)
E. coli Selection: Kanamycin (25 ug/mL)
Cell Selection: Neomycin
Fully Sequenced ORF: >MC212048 representing NM_178891
 Red=Cloning site Blue=ORF Orange=Stop codon

TTTTGTAATACGACTCACTATAGGGCGCCGGAATTCGTCGACTGGATCCGGTACCGAGGAGATCTGCC
 GCC**GCGATCGCC**

ATGTCGCTGAGCAAGAAAAGAAAGCTTGAGTCGGGGACAGCGGAGGCGCCGGCCGGAGGGGAGGGAG
 CCGAGGAGGAAAATGGCGGGGAGCAGGAGGCGGCCCCGCCACGACCCGGAGGACCAAGAGCGAGCGGGA
 CCAGCTGTACTACGAGTCTACTCCGACGTCTCGGTCCACGAGGAGATGATCGCCGACCAGTCCGCACC
 GAAGCCTACCGCTTAGGCATCCTGAAGAACTGGGCGCGCTGCGAGGCAAGACGGTGTGGACGTGGGCG
 CGGGCACCGCATTCTCAGCATTTCTGTGCCAGGCCGGGGCACGGCGCGTGTACGCGGTGGAGGCCAG
 CGCCATCTGGCAACAGGCCCGGGAGGTGGTGGCGCTCAACGGGTTGGAGGACCGCGTGCACGTCTGCCG
 GGCCCGGTGGAGACCGTGGAGCTGCCGGAGCGAGTGGACGCCATCGTCAGCGAGTGGATGGGCTACGGAC
 TTCTGCACGAGTCCATGCTGAGCTCCGTGCTCCACGCGCGGACCAATGGCTGAAGGAGGGCGGTCTCCT
 CCTGCCAGCTCCGCGGAGCTCTCGTGGCCCCGATTAGCGACCAGATGCTCGAGTGGCGTCTGGGTTTC
 TGGAGCCAGGTGAAGCAGCACTATGGCGTGGATATGAGCTGCATGGAGAGCTTCGCCACGCGCTGCCTCA
 TGGGCCATTCGGAGATCGTGGTGCAGGATCTGTCCGGAGAGGACGTGTGGCCCGGCCGACGCGCTTTGC
 CCAGCTCGAGCTGGCCCGAGCCGCGCTGGAGCAGGAGCTGGAGGCTGGTGTGGCGGGCGCTTCCGCTGC
 AGCTGCTATGTTCCGCGCCTCTACATGGTTTCGCGCTGTGTTTCAAGTGACCTTTCCCGAGGGGACT
 CGGAGAAACCTCTGGTGTGTCCACCTCGCCTTTTACCCGCGCCACCCACTGGAAGCAGGCGCTCCTCTA
 CTTGAACGAGCCGGTGGCGGTGGAACAAGATACGGACATTTCCGGAGAGATCACCTGCTGCCCTCCCCG
 GACAACCCCGCGCTCTGCGCATACTTCTGCGCTACAAAGTGGGAGACCATGAGGAAAAGACCAAAGACT
 TTGCCATGGAGGACTGA

ACGCGTACGCGGCCGCTCGAGCAGAAACTCATCTCAGAAGAGGATCTGGCAGCAAATGATATCCTGGATT
 ACAAGGATGACGACGATAAGGTTTAA

Restriction Sites: SgfI-MluI



[View online »](#)

ACCN:	NM_178891
Insert Size:	1137 bp
OTI Disclaimer:	Our molecular clone sequence data has been matched to the reference identifier above as a point of reference. Note that the complete sequence of our molecular clones may differ from the sequence published for this corresponding reference, e.g., by representing an alternative RNA splicing form or single nucleotide polymorphism (SNP).
Components:	The ORF clone is ion-exchange column purified and shipped in a 2D barcoded Matrix tube containing 10ug of transfection-ready, dried plasmid DNA (reconstitute with 100 ul of water).
Reconstitution Method:	<ol style="list-style-type: none">1. Centrifuge at 5,000xg for 5min.2. Carefully open the tube and add 100ul of sterile water to dissolve the DNA.3. Close the tube and incubate for 10 minutes at room temperature.4. Briefly vortex the tube and then do a quick spin (less than 5000xg) to concentrate the liquid at the bottom.5. Store the suspended plasmid at -20°C. The DNA is stable for at least one year from date of shipping when stored at -20°C.
RefSeq:	<u>NM_178891.5</u> , <u>NP_849222.3</u>
RefSeq Size:	2475 bp
RefSeq ORF:	1137 bp
Locus ID:	99890
UniProt ID:	<u>Q6Nzb1</u>
Cytogenetics:	3 F3

Gene Summary:

Arginine methyltransferase that can catalyze the formation of both omega-N monomethylarginine (MMA) and asymmetrical dimethylarginine (aDMA), with a strong preference for the formation of aDMA (PubMed:22904064, PubMed:26070566). Preferentially methylates arginyl residues present in a glycine and arginine-rich domain and displays preference for monomethylated substrates (By similarity). Specifically mediates the asymmetric dimethylation of histone H3 'Arg-2' to form H3R2me2a (By similarity). H3R2me2a represents a specific tag for epigenetic transcriptional repression and is mutually exclusive with methylation on histone H3 'Lys-4' (H3K4me2 and H3K4me3) (By similarity). Acts as a transcriptional repressor of various genes such as HOXA2, THBS1 and TP53 (PubMed:22904064). Repression of TP53 blocks cellular senescence (PubMed:22904064). Also methylates histone H2A and H4 'Arg-3' (H2AR3me and H4R3me, respectively). Acts as a regulator of DNA base excision during DNA repair by mediating the methylation of DNA polymerase beta (POLB), leading to the stimulation of its polymerase activity by enhancing DNA binding and processivity. Methylates HMGA1. Regulates alternative splicing events. Acts as a transcriptional coactivator of a number of steroid hormone receptors including ESR1, ESR2, PGR and NR3C1. Promotes fasting-induced transcriptional activation of the gluconeogenic program through methylation of the CRT2 transcription coactivator (PubMed:24570487). Methylates GPS2, protecting GPS2 from ubiquitination and degradation (PubMed:26070566). [UniProtKB/Swiss-Prot Function]

Transcript Variant: This variant (1) represents the shorter transcript, is intronless and encodes the functional protein.